

Datasheet for ABIN1413320

anti-CARTPT antibody (AA 31-116) (HRP)



Go to Product page

\sim				
()\	/e	r\ /	\cap	٨
()	<i>v</i> \leftarrow	ı vı	\Box	٧V

Quantity:	100 μL
Target:	CARTPT
Binding Specificity:	AA 31-116
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CARTPT antibody is conjugated to HRP
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CART	
Isotype:	IgG	
Predicted Reactivity:	Human,Mouse,Rat,Cow,Sheep,Pig	
Purification:	Purified by Protein A.	

Target Details

Target:	CARTPT
Alternative Name:	CART (CARTPT Products)
Background:	Synonyms: CART, CART prepropeptide, Cocaine and amphetamine regulated transcript,

CART_HUMAN.

Background: The CART gene encodes for a protein which has an important role in the regulation of appetite and body weight. The CART (cocaine- and amphetamine-regulated transcript) neuropeptide is an mRNA that changes in response to psychostimulant drug administration. Injection of CART peptides into the ventral tegmental area produces psychostimulant-like effects. CART localizes to areas of the central and peripheral nervous systems and is involved in feeding behavior when injected centrally. Expression of CART in the rat hypothalamus is modulated by nutritional status, and injection of synthetic CART peptide into the forebrain ventricular system suppresses food intake, indicating a possible role in hypothalamic control of energy homeostasis. Its identification in cell bodies and central terminals of vagal afferent neurons additionally suggests a role in brainstem mechanisms of meal termination and satiety.

Pathways:

Hormone Transport, Negative Regulation of Hormone Secretion, Carbohydrate Homeostasis, Feeding Behaviour

Application Details

Application Notes: IHC-P 1:200-400

IHC-F 1:100-500

Restrictions: For Research Use only

Handling

50 % Glycerol. Preservative: ProClin		
Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should handled by trained staff only. Handling Advice: Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase. Storage: -20 °C	Format:	Liquid
50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should handled by trained staff only. Handling Advice: Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase. Storage: -20 °C	Concentration:	1 μg/μL
Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should handled by trained staff only. Handling Advice: Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase. Storage: -20 °C	Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
handled by trained staff only. Handling Advice: Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase. Storage: -20 °C	Preservative:	ProClin
peroxidase. Storage: -20 °C	Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
	Handling Advice:	
Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Storage:	-20 °C
	Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

	l l	1:
-	เวทก	111111
	land	HHU

Expiry Date:

12 months