antibodies -online.com





anti-Taxilin beta antibody (AA 331-430) (HRP)



Go to	Prod	luct	page

()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL
Target:	Taxilin beta (TXLNB)
Binding Specificity:	AA 331-430
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Taxilin beta antibody is conjugated to HRP
Application:	Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Western Blotting (WB), Immunohistochemistry (Frozen Sections) (IHC (fro)), ELISA

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human TXLNB
Isotype:	IgG
Specificity:	This may have some cross-reactivity with TXLNA and TXLNG.
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Sheep,Pig,Horse
Purification:	Purified by Protein A.

Target Details

|--|--|--|

Target Details

Alternative Name:	TXLNB (TXLNB Products)
Background:	Synonyms: MDP77, LST001, C6orf198, dJ522B19.2, Beta-taxilin, Muscle-derived protein 77, hMDP77, TXLNB
	Background: Beta-taxilin promotes nerve regeneration and may be involved in intracellular
	vesicle transport. Expressed predominantly in cardiac and skeletal muscle, beta-taxilin binds to
	the coiled coil region of the syntaxin family members STX1A, STX3A, and STX4A. Beta- and
	gamma-taxilins, bind to the alpha subunit of the nascent polypeptide-associated complex
	(NAC) and affect its nuclear distribution, suggesting that the taxilin family is involved not only in
	the translational process through its interaction with NAC but also in the transcriptional process through its interaction with alphaNAC alone.
Gene ID:	167838
UniProt:	Q8N3L3
Application Details	
Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish
	peroxidase.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

	_	
-	II	1:
-	-222	una
-	Hand	1111()

Expiry Date:

12 months