Datasheet for ABIN1413935 anti-THO Complex 2 antibody (AA 101-200) (Cy5.5)

-online.com antibodies



Overview

Quantity:	100 µL	
Target:	THO Complex 2 (THOC2)	
Binding Specificity:	AA 101-200	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This THO Complex 2 antibody is conjugated to Cy5.5	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human THOC2	
lsotype:	lgG	
Cross-Reactivity:	Human	
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Pig,Horse,Rabbit	
Purification:	Purified by Protein A.	
Target Details		
Target:	THO Complex 2 (THOC2)	
Alternative Name:	Thoc2 (THOC2 Products)	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN1413935 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

Target Details

Background:	Synonyms: THO2, CXorf3, hTREX120, dJ506G2.1, THO complex subunit 2, THOC2		
	Background: Required for efficient export of polyadenylated RNA and spliced mRNA. Acts as		
	component of the THO subcomplex of the TREX complex which is thought to couple mRNA		
	transcription, processing and nuclear export, and which specifically associates with spliced		
	mRNA and not with unspliced pre-mRNA. TREX is recruited to spliced mRNAs by a		
	transcription-independent mechanism, binds to mRNA upstream of the exon-junction complex		
	(EJC) and is recruited in a splicing- and cap-dependent manner to a region near the 5' end of the		
	mRNA where it functions in mRNA export to the cytoplasm via the TAP/NFX1 pathway. The		
	TREX complex is essential for the export of Kaposi's sarcoma-associated herpesvirus (KSHV)		
	intronless mRNAs and infectious virus production. THOC2 (and probably the THO complex) is		
	involved in releasing mRNA from nuclear speckle domains. Required for NXF1 localization to		
	the nuclear rim.		
Gene ID:	57187		
UniProt:	Q8NI27		

Application Details

Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN1413935 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

1.1	(1:
Н	land	ling
		3

Expiry Date:

12 months

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN1413935 | 03/07/2024 | Copyright antibodies-online. All rights reserved.