

Datasheet for ABIN1414650

anti-RUNX3 antibody (pSer338) (Cy5.5)



Overview

Quantity:	100 μL
Target:	RUNX3
Binding Specificity:	pSer338
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RUNX3 antibody is conjugated to Cy5.5
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)),
	Immunofluorescence (Cultured Cells) (IF (cc))
Product Details	
Immunogen:	KLH conjugated synthetic phosphopeptide derived from human RUNX3 around the
	phosphorylation site of Ser338
Isotype:	IgG
Predicted Reactivity:	Human
Purification:	Purified by Protein A.
Target Dataila	
Target Details	
Target:	RUNX3
Alternative Name:	RUNX3 (RUNX3 Products)

Target Details

Storage Comment:

factor, alpha 3 subunit, CBF-alpha 3, Acute myeloid leukemia 2 protein, OncogeneAML-2, Polyomavirus enhancer-binding protein 2 alpha C subunit, PEBP2-alpha C, PEA2-alpha C, SL3-3 enhancer factor 1 alpha C subunit, SL3/AKV core-binding factor alpha C subunit, Core-binding factor, alpha 3 subunit, CBF-alpha 3, Acute myeloid leukemia 2 protein, Oncogene AMI-2, Polyomavirus enhancer-binding protein 2 alpha C subunit, PEBP2-alpha C, RUNX3.HUMAN, PEA2-alpha C, SL3-3 enhancer factor 1 alpha C subunit, SL3/AKV core-binding factor alpha C subunit. Background: RUNX3 binds to the core site of murine Leukemia virus, the core sequences in the enhancer of the polyomavirus, and also to the enhancers of the T-cell receptor genes. May be involved in the control of cellular proliferation and/or differentiation (By similarity), Heterodimer of an alpha and a beta subunit. The alpha subunit binds DNA as a monomer and through the Runt domain.DNA-binding is increased by heterodimerization. Interacts with TLE1 and SUV39H1, Subcellular location in Nucleus. Gene ID: 864 Application Details IF(IHC-P) 1:50-200 IF(ICC) 1:50-200 IF(ICC) 1:50-200 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. Preservative: ProClin	l arget Details	
SUV39H1, Subcellular location in Nucleus. Gene ID: 864 Application Details Application Notes: IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	Background:	Polyomavirus enhancer-binding protein 2 alpha C subunit, PEBP2-alpha C, PEA2-alpha C, SL3-3 enhancer factor 1 alpha C subunit, SL3/AKV core-binding factor alpha C subunit, Core-binding factor, alpha 3 subunit, CBF-alpha 3, Acute myeloid leukemia 2 protein, Oncogene AML-2, Polyomavirus enhancer-binding protein 2 alpha C subunit, PEBP2-alpha C, RUNX3_HUMAN, PEA2-alpha C, SL3-3 enhancer factor 1 alpha C subunit, SL3/AKV core-binding factor alpha C subunit. Background: RUNX3 binds to the core site of murine Leukemia virus, the core sequences in the enhancer of the polyomavirus, and also to the enhancers of the T-cell receptor genes. May be involved in the control of cellular proliferation and/or differentiation (By similarity). Heterodimer
Gene ID: 864 Application Details IF(IHC-P) 1:50-200 Application Notes: IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.		· · · · · · · · · · · · · · · · · · ·
Application Details Application Notes: IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.		
Application Notes: IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 Restrictions: For Research Use only Handling Format: Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	Gene ID:	864
IF(IHC-F) 1:50-200 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	Application Details	
IF(ICC) 1:50-200 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	Application Notes:	IF(IHC-P) 1:50-200
Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.		IF(IHC-F) 1:50-200
Handling Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.		IF(ICC) 1:50-200
Format: Liquid Concentration: 1 μg/μL Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. Preservative: ProClin This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	Restrictions:	For Research Use only
Concentration: 1 μg/μL Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. Preservative: ProClin This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	Handling	
Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. Preservative: ProClin This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	Format:	Liquid
50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	Concentration:	1 μg/μL
Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
handled by trained staff only.	Preservative:	ProClin
Storage: -20 °C	Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
	Storage:	-20 °C

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

ш	Jand	lina
г	land	1111()
•	10110	9

Expiry Date:

12 months