

[Go to Product page](#)

Datasheet for ABIN1414729

**anti-Complement C3b antibody (Cy7)**

## Overview

Quantity:	100 µL
Target:	Complement C3b (C3b)
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Complement C3b antibody is conjugated to Cy7
Application:	Western Blotting (WB)

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Complement C3 beta chain
Isotype:	IgG
Specificity:	This antibody could recognize C3-beta-c
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Dog,Cow,Pig,Horse,Guinea Pig
Purification:	Purified by Protein A.

## Target Details

Target:	Complement C3b (C3b)
Alternative Name:	Complement C3 beta Chain ( <a href="#">C3b Products</a> )
Background:	Synonyms: ASP, C3a, C3b, AHUS5, ARMD9, CPAMD1, HEL-S-62p, Complement C3, C3 and PZP-

## Target Details

like alpha-2-macroglobulin domain-containing protein 1, C3, CO3\_HUMAN

Background: C3 plays a central role in the activation of the complement system. Its processing by C3 convertase is the central reaction in both classical and alternative complement pathways. After activation C3b can bind covalently, via its reactive thioester, to cell surface carbohydrates or immune aggregates. Derived from proteolytic degradation of complement C3, C3a anaphylatoxin is a mediator of local inflammatory process. In chronic inflammation, acts as a chemoattractant for neutrophils (By similarity). It induces the contraction of smooth muscle, increases vascular permeability and causes histamine release from mast cells and basophilic leukocytes. C3-beta-c: Acts as a chemoattractant for neutrophils in chronic inflammation. Acylation stimulating protein: adipogenic hormone that stimulates triglyceride (TG) synthesis and glucose transport in adipocytes, regulating fat storage and playing a role in postprandial TG clearance. Appears to stimulate TG synthesis via activation of the PLC, MAPK and AKT signaling pathways. Ligand for C5AR2. Promotes the phosphorylation, ARRB2-mediated internalization and recycling of C5AR2 (PubMed:8376604, PubMed:2909530, PubMed:9059512, PubMed:10432298, PubMed:15833747, PubMed:16333141, PubMed:19615750).

Gene ID:	718
UniProt:	<a href="#">P01024</a>
Pathways:	<a href="#">Complement System</a> , <a href="#">Regulation of Leukocyte Mediated Immunity</a> , <a href="#">Positive Regulation of Immune Effector Process</a> , <a href="#">Regulation of G-Protein Coupled Receptor Protein Signaling</a>

## Application Details

Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE, which

Handling

---

	should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months