

Datasheet for ABIN1414911 anti-ENPP1 antibody (AA 41-140) (HRP)



()	11	OF	· \ /	-	1 A /
	11/	-	1//	-	1/1/
	v	er	v		v v

Overview	
Quantity:	100 μL
Target:	ENPP1
Binding Specificity:	AA 41-140
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ENPP1 antibody is conjugated to HRP
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human ENPP1
Isotype:	IgG

Immunogen:	KLH conjugated synthetic peptide derived from human ENPP1
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	ENPP1
Alternative Name:	ENPP1 (ENPP1 Products)

Target Details

Background:	Synonyms: M6S1, NPP1, NPPS, PC-1, PCA1, ARHR2, COLED, PDNP1, Ectonucleotide
	pyrophosphatase/phosphodiesterase family member 1, E-NPP 1, Membrane component
	chromosome 6 surface marker 1, Phosphodiesterase I/nucleotide pyrophosphatase 1, Plasma-
	cell membrane glycoprotein PC-1, ENPP1, PC1
	Background: By generating PPi, plays a role in regulating pyrophosphate levels, and functions in
	bone mineralization and soft tissue calcification. PPi inhibits mineralization by binding to
	nascent hydroxyapatite (HA) crystals, thereby preventing further growth of these crystals.
	Preferentially hydrolyzes ATP, but can also hydrolyze other nucleoside 5' triphosphates such as
	GTP, CTP, TTP and UTP to their corresponding monophosphates with release of
	pyrophosphate and diadenosine polyphosphates, and also 3',5'-cAMP to AMP. May also be
	involved in the regulation of the availability of nucleotide sugars in the endoplasmic reticulum
	and Golgi, and the regulation of purinergic signaling. Appears to modulate insulin sensitivity and
	function.
Gene ID:	5167
UniProt:	P22413
Pathways:	Regulation of Carbohydrate Metabolic Process
Application Details	
Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.

Handling

	peroxidase.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months