antibodies -online.com





anti-Chromosome 5 Open Reading Frame 39 (C5orf39) (AA 50-100) antibody (Cy5.5)



Go to Product page

()	۱ ۱	\cap	r	/1	\cap	۱ ۸	1
0	'V	ㄷ	ı١	νı	\Box	V	۷

Quantity:	100 μL	
Target:	Chromosome 5 Open Reading Frame 39 (C5orf39)	
Binding Specificity:	AA 50-100	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	Cy5.5	
Application:	Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Annexin-2 receptor	
Isotype:	IgG	
Predicted Reactivity:	Human	
Purification:	Purified by Protein A.	

Target Details

Target:	Chromosome 5 Open Reading Frame 39 (C5orf39)
Alternative Name:	AX2R (C5orf39 Products)
Background: Synonyms: LOC389289, ANXA2R, Annexin 2 receptor, Annexin II receptor, AX2R	

Chromosome 5 open reading frame 39, AX2R_HUMAN.

Background: ANXA2R (annexin-2 receptor), also known as AX2R or C5orf39, is a 193 amino acid protein that is widely expressed and may act as an annexin II receptor on marrow stromal cells to induce osteoclast formation. In addition, ANXA2R is highly expressed in lymphocytes and is also found in resting CD4+ and CD8+ T cells. The gene encoding ANXA2R maps to human chromosome 5, which contains 181 million base pairs and comprises nearly 6 % of the human genome. Chromosome 5 is associated with Cockayne syndrome through the ERCC8 gene and familial adenomatous polyposis through the adenomatous polyposis coli (APC) tumor suppressor gene. Treacher Collins syndrome is also chromosome 5-associated and is caused by insertions or deletions within the TCOF1 gene. Deletion of the p arm of chromosome 5 leads to Cri du chat syndrome, while deletion of the q arm or of chromosome 5 altogether is common in therapy-related acute myelogenous leukemias and myelodysplastic syndrome.

Application Details

Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months