

Datasheet for ABIN1416436

anti-UBE2G2 antibody (AA 100-165) (HRP)



_				
()	1//	rv	IO	Λ/
()	VC	. I V	1	v v

Quantity:	100 μL
Target:	UBE2G2
Binding Specificity:	AA 100-165
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This UBE2G2 antibody is conjugated to HRP
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Ube2G2/UBC7
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Cow, Pig, Horse, Chicken, Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	UBE2G2
Alternative Name:	Ube2G2 (UBE2G2 Products)
Background: Synonyms: Human ubiquitin conjugating enzyme G29, UB2G2_HUMAN, UBC 7, UBC7, UBC7	

homolog yeast, UBE2G2, Ubiquitin carrier protein G2, Ubiquitin conjugating enzyme 7, Ubiquitin conjugating enzyme E2 G2, Ubiquitin conjugating enzyme E2G 2 homologous to yeast UBC7, Ubiquitin conjugating enzyme E2G 2 UBC7 homolog yeast, Ubiquitin conjugating enzyme E2G 2, Ubiquitin conjugating enzyme G2, Ubiquitin protein ligase G2, Ubiquitin-conjugating enzyme E2 G2, Ubiquitin-protein ligase G2.

Background: UBE2G2 (Ubiquitin-conjugating enzyme E2 G2), also known as UBC7, is a 165 amino acid protein involved in ubiquitin-mediated protein degradation. Ubiquitination is an important mechanism through which three classes of enzymes act in concert to target short-lived or abnormal proteins for destruction. The three classes of enzymes involved in ubiquitination are the ubiquitin-activating enzymes (E1s), the ubiquitin-conjugating enzymes (E2s) and the ubiquitin-protein ligases (E3s). UBE2G2 is an E2 ubiquitin-conjugating enzyme that acts to catalyze the covalent attachment of ubiquitins to various proteins. Expressed throughout the body, UBE2G2 shares 100 % sequence identity with its mouse counterpart and is thought to be involved in endoplasmic reticulum-associated degradation (ERAD). Two isoforms of UBE2G2 exist due to alternative splicing events.

Gene ID:

7327

IHC-F 1:100-500

Application Details

Application Notes:	IHC-P 1:200-400

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.

Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months