# antibodies .- online.com





Datasheet for ABIN1416557

# anti-UBE2H antibody (AA 101-183) (Cy3)

Go to Product page

_			
	N/0	r\/I	ew
$\sim$	' v C	1 V I	C V V

Quantity:	100 μL
Target:	UBE2H
Binding Specificity:	AA 101-183
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This UBE2H antibody is conjugated to Cy3
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human Ube2H/UBCH2
Isotype:	IgG
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Chicken,Rabbit
Purification:	Purified by Protein A.

# **Target Details**

Target:	UBE2H
Alternative Name:	Ube2H (UBE2H Products)
Background:	Synonyms: E2 20K, Ubc H2, UBC8, UBCH 2, UBCH, UBCH2, Ube2h, UBE2H_HUMAN, Ubiquitin

carrier protein H, Ubiquitin conjugating enzyme E2 H, Ubiquitin conjugating enzyme E2H UBC8 homolog yeast, Ubiquitin conjugating enzyme E2H, Ubiquitin protein ligase H, Ubiquitin-conjugating enzyme E2 H, Ubiquitin-conjugating enzyme E2-20K, Ubiquitin-protein ligase H. Background: UBE2H, also known as UBC8, UBCH, UBCH2 or E2-20K, is a 183 amino acid protein involved in ubiquitin-mediated protein degradation. Ubiquitination is an important mechanism through which three classes of enzymes act in concert to target short-lived or abnormal proteins for destruction. The three classes of enzymes involved in ubiquitination are the ubiquitin-activating enzymes (E1s), the ubiquitin-conjugating enzymes (E2s) and the ubiquitin-protein ligases (E3s). One of several members of the ubiquitin-conjugating enzyme family, UBE2H functions as an E2 ubiquitin-conjugating enzyme that acts to catalyze the covalent attachment of ubiquitin residues to various proteins, including Histone H2A. UBE2H shares 100 % identity with its mouse counterpart and 98 % identity with its frog and zebrafish homologs, suggesting a conserved function between species. Multiple isoforms of UBE2H exist due to alternative splicing events.

Gene ID:

7328

#### **Application Details**

Application	Notes:

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

## Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

		1.	
$\vdash$	land	lın	
	ıaııu	1111	U

Expiry Date:

12 months