antibodies .- online.com





anti-GLIPR2 antibody (AA 51-154) (Cy3)



		Go to Product page

()	1 /	0	rv	/ 1 /	71	Α.
	1//	\vdash	1 \/	16		1/1/
\sim	v	\sim	1 V	١,	_	v v

Quantity:	100 μL
Target:	GLIPR2
Binding Specificity:	AA 51-154
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GLIPR2 antibody is conjugated to Cy3
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human GLIPR2
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Sheep, Pig, Horse, Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	GLIPR2
Alternative Name:	GLIPR2 (GLIPR2 Products)
Background:	Synonyms: GAPR 1, GAPR-1, GAPR1_HUMAN, GLI pathogenesis related 2, Glioma pathogenesis

related protein 2, Glioma pathogenesis-related protein 2, GliPR 2, GLIPR2, Golgi associated plant pathogenesis related protein 1, Golgi associated PR 1 protein, Golgi-associated plant pathogenesis-related protein 1, Golgi-associated PR-1 protein, OTTMUSP00000007558, RP11-421H8.5, RP23-209M8.2, C77180, 5730414A08Rik, C9orf19.

Background: Cysteine-rich secretory proteins (CRISPs) represent a family of evolutionarily conserved proteins that may play a role in the innate immune system and are transcriptionally regulated by androgens in several tissues. GAPR-1 (Golgi-associated plant pathogenesis-related protein 1), also known as GLIPR2, is a 154 amino acid lipid anchor protein belonging to the CRISP family. GAPR-1 also shares similarity with the pathogenesis-related protein (PR) superfamily, and may play an important role in the immune system. Existing as a homodimer, GAPR-1 is highly expressed in lung and peripheral leukocytes with minor expression in liver and kidney. Containing a conserved sperm-coating protein (SCP) domain, GAPR-1 binds to negatively charged lipids and may be involved in the differentiation of epithelial cells into mesenchymal cells. Increased expression of GAPR-1 in kidney may contribute to the development of fibrosis.

Application Details

Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

		1.	
\vdash	land	ling	
1 1	ıaııu	11111	۰

Expiry Date:

12 months