Datasheet for ABIN1417087 anti-FNDC1 antibody (AA 21-120) (Cy5.5)

antibodies .-online.com



$\sim$			
()	VP	rvi	ew
$\sim$	v C		C * *

Quantity:	100 µL	
Target:	FNDC1	
Binding Specificity:	AA 21-120	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This FNDC1 antibody is conjugated to Cy5.5	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human FNDC1	
Isotype:	IgG	
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Horse,Rabbit	
Purification:	Purified by Protein A.	

## Target Details

Target:	FNDC1
Alternative Name:	FNDC1 (FNDC1 Products)
Background:	Synonyms: Activation-associated cDNA protein, AGS8, Expressed in synovial lining protein,

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN1417087 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

	fibronectin type III domain containing 2, Fibronectin type III domain-containing protein 1,	
	FNDC1, FNDC1_HUMAN, FNDC2, MEL4B3.	
	Background: FNDC1, also known as MEL4B3, Ags8, Expressed in synovial lining protein and	
	Activation-associated cDNA protein, is a 1888 amino acid secreted protein that contains five	
	fibronectin type-III domains. FNDC1 is moderately expressed in skeletal muscle, pancreas,	
	heart, kidney, spinal cord, ovary and lung. Expression of FNDC1 is induced in response to	
	hypoxia in ventricular cardiomyocytes. Since FNDC1 interacts with G and G©, it is likely that	
	FNDC1 is an activator for G-protein signaling. Though normally absent in healthy skin, FNDC1	
	expression is induced by TGF-beta signaling in skin tumors and psoriasis. There are two	
	isoforms of FNDC1 that are produced as a result of alternative splicing events.	
Gene ID:	84624	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/2 | Product datasheet for ABIN1417087 | 03/07/2024 | Copyright antibodies-online. All rights reserved.