antibodies .- online.com







anti-RENBP antibody (AA 101-200) (Cy3)



oo to i loadot pago

Overview
Quantity:

100 μL

Target: RENBP

Binding Specificity: AA 101-200

Reactivity: Human, Mouse

Host: Rabbit

Clonality: Polyclonal

Conjugate: This RENBP antibody is conjugated to Cy3

Application: Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded

Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human RENBP
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Cow,Sheep,Horse
Purification:	Purified by Protein A.

Target Details

Target:	RENBP
Alternative Name:	RENBP (RENBP Products)

Target Details

Background:	Synonyms: AGE, EC 5.1.3.8, GlcNAc 2 epimerase, GlcNAc 2-epimerase, N acetyl D glucosamine
	2 epimerase, N-acetyl-D-glucosamine 2-epimerase, N-acylglucosamine 2-epimerase, RBP,
	RENBP, RENBP_HUMAN, Renin binding protein, Renin-binding protein, RNBP.
	Background: RENBP is a 427 amino acid protein that is a proteinaceous renin inhibitor. In
	inhibiting renin, RENBP forms a complex with it, a high molecular weight renin. RENBP contains
	a leucine zipper domain, which is essential for its dimerization with renin. RENBP can catalyze
	the interconversion of N-acetylglucosamine to N-acetylmannosamine, indicating that it is a
	GlcNAc 2-epimerase. Sequences of porcine, human and rat renin-binding proteins are highly
	homologous. The RENBP gene is conserved in dog, mouse, rat and zebrafish, and maps to
	human chromosome Xq28 between DXS52 and G6PD. Rat Renbp is located on chromosome X
	at Xq37 close to marker DXWox3 and falls outside the BP QTL regions on chromosome X.
Gene ID:	5973
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months