

Datasheet for ABIN1417310

anti-OSTM1 antibody (AA 21-120) (Cy7)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	OSTM1
Binding Specificity:	AA 21-120
Reactivity:	Human, Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This OSTM1 antibody is conjugated to Cy7
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human OSTM1
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Pig
Purification:	Purified by Protein A.

Target Details

Target:	OSTM1
Alternative Name:	OSTM1 (OSTM1 Products)

Target Details

Background:	Synonyms: GL, GIPN, OPTB5, HSPC019, Osteopetrosis-associated transmembrane protein 1, Chloride channel 7 beta subunit, OSTM1, UNQ6098/PRO21201 Background: OSTM1 (osteopetrosis associated transmembrane protein 1), also known as gl (gray-lethal) or HSPC019, is a 338 amino acid single-pass type I membrane protein that is expressed primarily in osteoclasts and melanocytes as well as brain, kidney and spleen. Bone autosomal recessive osteopetrosis (ARO) is the most severe form of hereditary bone disease whose cellular basis is in the osteoclast and is characterized by abnormally dense bone, due to defective resorption of immature bone. ARO is suggested to be caused by mutations in the OSTM1 gene. The disorder occurs in two forms: a severe autosomal recessive form occurring in utero, infancy, or childhood, and a benign autosomal dominant form occurring in adolescence or adulthood. Defects in the OSTM1 gene are also the cause of the spontaneous gl mutant, which is responsible for a coat color defect in mice.
Gene ID:	28962
UniProt:	Q86WC4

Application Details

Application Notes:	FCM 1:20-100 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Handling

Expiry Date: 12 months