

Datasheet for ABIN1417810

anti-C9ORF72 antibody (AA 391-481) (HRP)



Go to Product page

_		erview/				
	1//	r	1//	\triangle	۸/	
	V		VI		/ V	

Alternative Name:

Quantity:	100 μL	
Target:	C90RF72	
Binding Specificity:	AA 391-481	
Reactivity:	Human, Rat, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This C9ORF72 antibody is conjugated to HRP	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)),	
	Immunohistochemistry (Frozen Sections) (IHC (fro))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human C9orf72	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Predicted Reactivity:	Dog,Cow,Pig,Horse,Chicken	
Purification:	Purified by Protein A.	
Target Details		
Target:	C90RF72	

C9orf72 (C9ORF72 Products)

Target Details

Synonyms: ALSFTD, FTDALS, Protein C9orf72, C9orf72 Background: Chromosome 9 consists of about 145 million bases and 4 % of the human genome and encodes nearly 900 genes. Considered to play a role in gender determination, deletion of the distal portion of 9p can lead to development of male to female sex reversal, the phenotype of a female with a male X,Y genotype. Hereditary hemorrhagic telangiectasia, which is characterized by harmful vascular defects, is associated with the chromosome 9 gene encoding endoglin protein, ENG. Familial dysautonomia is also associated with chromosome 9 though through the gene IKBKAP. Notably, chromosome 9 encompasses the largest interferon family gene cluster. Chromosome 9 is partnered with chromosome 22 in the translocation	
genome and encodes nearly 900 genes. Considered to play a role in gender determination, deletion of the distal portion of 9p can lead to development of male to female sex reversal, the phenotype of a female with a male X,Y genotype. Hereditary hemorrhagic telangiectasia, which is characterized by harmful vascular defects, is associated with the chromosome 9 gene encoding endoglin protein, ENG. Familial dysautonomia is also associated with chromosome 9 though through the gene IKBKAP. Notably, chromosome 9 encompasses the largest interferon family gene cluster. Chromosome 9 is partnered with chromosome 22 in the translocation	
deletion of the distal portion of 9p can lead to development of male to female sex reversal, the phenotype of a female with a male X,Y genotype. Hereditary hemorrhagic telangiectasia, which is characterized by harmful vascular defects, is associated with the chromosome 9 gene encoding endoglin protein, ENG. Familial dysautonomia is also associated with chromosome 9 though through the gene IKBKAP. Notably, chromosome 9 encompasses the largest interferon family gene cluster. Chromosome 9 is partnered with chromosome 22 in the translocation	
phenotype of a female with a male X,Y genotype. Hereditary hemorrhagic telangiectasia, which is characterized by harmful vascular defects, is associated with the chromosome 9 gene encoding endoglin protein, ENG. Familial dysautonomia is also associated with chromosome 9 though through the gene IKBKAP. Notably, chromosome 9 encompasses the largest interferon family gene cluster. Chromosome 9 is partnered with chromosome 22 in the translocation	
is characterized by harmful vascular defects, is associated with the chromosome 9 gene encoding endoglin protein, ENG. Familial dysautonomia is also associated with chromosome 9 though through the gene IKBKAP. Notably, chromosome 9 encompasses the largest interferon family gene cluster. Chromosome 9 is partnered with chromosome 22 in the translocation	
encoding endoglin protein, ENG. Familial dysautonomia is also associated with chromosome 9 though through the gene IKBKAP. Notably, chromosome 9 encompasses the largest interferon family gene cluster. Chromosome 9 is partnered with chromosome 22 in the translocation	
though through the gene IKBKAP. Notably, chromosome 9 encompasses the largest interferon family gene cluster. Chromosome 9 is partnered with chromosome 22 in the translocation	
family gene cluster. Chromosome 9 is partnered with chromosome 22 in the translocation	
leading to the aberrant production of BCR-ABL fusion protein often found in leukemias. The	
C9orf72 gene product has been provisionally designated C9orf72 pending further	
characterization. There are two isoforms of C9orf72 that are produced as a result of alternative	
splicing events.	
203228	
Q96LT7	
WB 1:300-5000	
IHC-P 1:200-400	
IHC-F 1:100-500	
For Research Use only	
Liquid	
1 μg/μL	
Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
50 % Glycerol.	
ProClin	
This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.	
•	

Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months