antibodies .- online.com





Datasheet for ABIN1418256

anti-IQCG antibody (Cy5)

Overview	
Quantity:	100 μL
Target:	IQCG
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IQCG antibody is conjugated to Cy5
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	

Immunogen:	KLH conjugated synthetic peptide derived from human IQCG
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

Target Details	
Target:	IQCG
Alternative Name:	IQCG (IQCG Products)
Background:	Synonyms: IQ domain containing protein G, IQ mot containing G, DKFZp434B227, FLJ11667, FLJ23571, FLJ37775, IQCG_HUMAN.
	Background: IQCG is a 443 amino acid protein containing one IQ domain. Widely distributed in
	nature, the IQ domain forms an amphiphilic seven-turn -helix capable of binding calmodulin in a

Target Details

Ca2+-independent manner. The level of intracellular calcium is tightly regulated in all eukaryotic cells. A modest increase in this level can result in a myriad of physiological responses, most of which are mediated by calmodulin (CaM), the universal calcium sensor. In acute T-lymphoid/myeloid leukemia, IQCG forms a complex with Nup98, an O-linked glycoprotein and a component of the nuclear pore complex. NUP98-IQCG complex bind co-activators and/or co-repressors, which suggest a role in transcriptional regulation.Nup98-IQCG complex inhibits 32Dcl3 cell apoptosis induced by Arabinofuranosylcytosine (Ara-C) and partially blocks granulocyte differentiation induced by G-CSF. IQCG exists as two isoforms due to alternatively splicing events.

Gene ID:

Storage:

Expiry Date:

Storage Comment:

84223

Application Details

Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

handled by trained staff only.

-20 °C

12 months