antibodies -online.com





anti-CINP antibody (Cy5)



_							
0	V	е	r١	/	е	٧	V

- Overview		
Quantity:	100 μL	
Target:	CINP	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CINP antibody is conjugated to Cy5	
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human CINP	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Purification:	Purified by Protein A.	
Target Details		
Target:	CINP	
Alternative Name:	CINP (CINP Products)	
Background:	Synonyms: CINP protein, CDK2-interacting protein, CINP, CINP_HUMAN, Cyclin dependent kinase 2 interacting protein, Cyclin-dependent kinase 2-interacting protein, MGC849. Background: Cell cycle progression is controlled in part by a family of cyclin proteins and cyclin	

dependent kinases (Cdks). Cdk proteins work in concert with the cyclins to phosphorylate key

Target Details

substrates involved in each phase of cell cycle progression. Specifically, Cdk2 interacts with Cyclins A, B1, B3, D, or E to control cell cycle progression. The Cyclin-dependent kinase 2-interacting protein (CINP) interacts with components of the replication complex and Cdk2 and Cdc7, thereby providing a functional and physical link between Cdk2 and Cdc7 during firing of the origins of replication. However, CINP is phopshorylated by Cdc7, but not by Cdk2. CINP also interacts with ATR-interacting protein and regulates ATR-dependent signaling, resistance to replication stress and G2 checkpoint integrity.

Gene ID:

Storage:

Expiry Date:

Storage Comment:

51550

-20 °C

12 months

Application Details

Application Notes:	IF(IHC-P) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should handled by trained staff only.	

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.