



[Go to Product page](#)

Datasheet for ABIN1418752
anti-POP1 antibody (HRP)

Overview

Quantity:	100 µL
Target:	POP1
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This POP1 antibody is conjugated to HRP
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human ASC2/POP1
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

Target Details

Target:	POP1
Alternative Name:	POP1 (POP1 Products)
Background:	Synonyms: PAAD domain only protein, POP1, PYDC1, Pyrin domain containing 1, Pyrin only protein 1, PYDC1_HUMAN. Background: The death domain (DD) superfamily of proteins share one or more of the following domains: the DD, DED (death-effector domain), CARD (caspase-recruitment domain) and PYD

Target Details

(Pyrin domain). Each of these domains is characterized by a canonical death domain fold, which consists of a bundle of five or six antiparallel α -helices. As their names suggest, these domains play prominent roles in programmed cell death. ASC2 (apoptosis-associated speck-like protein containing a CARD 2), also known as Pyrin-only protein 1 or PADD-only protein 1, is an 89 amino acid member of the DD superfamily that contains one Pyrin domain. Localized to the cytoplasm, ASC2 interacts with ASC to modulate NF- κ B and pro-caspase-1 regulation. ASC2 is predominantly expressed in monocytes, macrophages and granulocytes.

Gene ID: 10940

Application Details

Application Notes: WB 1:300-5000
IHC-P 1:200-400

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 μ g/ μ L

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling Advice: Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.

Storage: -20 $^{\circ}$ C

Storage Comment: Store at -20 $^{\circ}$ C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months