.-online.com antibodies

## Datasheet for ABIN1418795 anti-BAGE4 antibody (Cy3)



| Overview          |  |
|-------------------|--|
| Quantity:         | 100 µL   |
| Target:           | BAGE4  |
| Reactivity:       | Human  |
| Host:             | Rabbit   |
| Clonality:        | Polyclonal   |
| Conjugate:        | This BAGE4 antibody is conjugated to Cy3   |
| Application:      | Immunofluorescence (Paraffin-embedded Sections) (IF (p))   |
| Product Details   |  |
| Immunogen:        | KLH conjugated synthetic peptide derived from human BAGE4  |
| Isotype:          | IgG  |
| Cross-Reactivity: | Human  |
| Purification:     | Purified by Protein A.   |
| Target Details    |  |
| Target:           | BAGE4  |
| Alternative Name: | BAGE4 (BAGE4 Products)   |
| Background:       | Synonyms: B melanoma antigen 4, B melanoma antigen family, member 4, BAGE4,<br>BAGE4_HUMAN, Cancer/testis antigen 2.4, CT2.4, MGC119851, MGC119852, MGC119853,<br>MLL3P. |

Background: Members of the BAGE gene family encode antigens that are recognized by

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN1418795 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

## Target Details

|                     | cytotoxic T lymphocytes and are also known as CT (cancer/testis) antigens. Generated by         |
|---------------------|---|
|                     | juxtacentromeric shuffling of the MLL3 gene, the ancestral BAGE gene was expanded by            |
|                     | acrocentric exchanges and/or juxtacentromeric movements. BAGE, also known as B                  |
|                     | melanoma antigen 1 or cancer/testis antigen 2.1, is a 43 amino acid member of the BAGE          |
|                     | family. Generally, BAGE proteins are not expressed in normal tissues, except in testis, but are |
|                     | expressed highly in melanomas, bladder carcinomas, head and neck squamous cell                  |
|                     | carcinomas, and lung and breast cancer carcinomas. BAGE proteins are not expressed in renal,    |
|                     | colorectal and prostatic carcinomas, leukemias and lymphomas.                                   |
| Gene ID:            | 3624  |
| Application Details |   |

| Application Notes: | IF(IHC-P) 1:50-200    |
|--------------------|-----------------------|
| Restrictions:      | For Research Use only |

## Handling

| Format:            | Liquid   |
|--------------------|--|
| Concentration:     | 1 µg/µL  |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.        |
| Preservative:      | ProClin  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage:           | -20 °C   |
| Storage Comment:   | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                                  |
| Expiry Date:       | 12 months  |