.-online.com antibodies

Datasheet for ABIN1418832 anti-COPS4 antibody (Cy5)



Overview

Overview	
Quantity:	100 µL
Target:	COPS4
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This COPS4 antibody is conjugated to Cy5
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human COPS4
lsotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.
Target Details	
Target:	COPS4
Alternative Name:	COPS4 (COPS4 Products)
Background:	Synonyms: CH4, COP9 constitutive photomorphogenic homolog subunit 4, COP9 complex homolog subunit 4, COP9 complex S4, COP9 constitutive photomorphogenic homolog subunit 4, COP9 signalosome complex subunit 4, COP9 signalosome subunit 4, COPS4, CSN complex

subunit 4, CSN4, CSN4 SCHPO, CSN4_HUMAN, DCH4, JAB1 containing signalosome subunit 4,

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN1418832 | 03/06/2024 | Copyright antibodies-online. All rights reserved.

	JAB1-containing signalosome subunit 4, MGC94896, SGN4, Signalosome subunit 4. Background: Component of the COP9 signalosome complex (CSN), a complex involved in various cellular and developmental processes. The CSN complex is an essential regulator of the ubiquitin (UbI) conjugation pathway by mediating the deneddylation of the cullin subunits of SCF-type E3 ligase complexes, leading to decrease the UbI ligase activity of SCF-type complexes such as SCF, CSA or DDB2. The complex is also involved in phosphorylation of p53/TP53, c-jun/JUN, IkappaBalpha/NFKBIA, ITPK1 and IRF8/ICSBP, possibly via its association with CK2 and PKD kinases. CSN-dependent phosphorylation of TP53 and JUN promotes and protects degradation by the UbI system, respectively.
Gene ID:	51138
Pathways:	Cell Division Cycle
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months