antibodies -online.com





anti-SH3RF1 antibody (Cy3)



Go to Product page

\sim			
	N/P	r\/I	i⊢₩

Background:

Quantity:	100 μL	
Target:	SH3RF1	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This SH3RF1 antibody is conjugated to Cy3	
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human SH3MD2/RNF142	
Immunogen:	KLH conjugated synthetic peptide derived from human SH3MD2/RNF142	
Isotype:	IgG	
Isotype: Cross-Reactivity:	IgG Human, Mouse, Rat	
Isotype: Cross-Reactivity: Purification:	IgG Human, Mouse, Rat	
Isotype: Cross-Reactivity: Purification: Target Details	IgG Human, Mouse, Rat Purified by Protein A.	

SH3MD2, SH3 multiple domains 2, SH3RF1.

Synonyms: Plenty of SH3 domains, Plenty of SH3s, POSH, Ring finger protein 142, RNF 142,

RNF142, SH3 domain containing ring finger 1, SH3 domain containing RING finger protein 1,

Background: SH3MD2 acts as a scaffold protein, contributes to Rac-induced signal

Target Details

transduction such as JNKs (MAPK8 and MAPK9) activation and induces apoptosis. Within a signaling complex, it probably recruits protein kinases such as MAP3K10 or MAP3K11 which are in turn activated leading to the sequential activation of MAP2K4, MAP2K7 and JNKs (MAPK8 and MAPK9). SH3MD2 may be involved in targeting of HIV-1 GAG and GAG-POL polyproteins to the plasma membrane. This gene encodes a protein containing an N-terminus RING-finger, four SH3 domains, and a region implicated in binding of the Rho GTPase Rac. Via the RING-finger, the encoded protein has been shown to function as an ubiquitin-protein ligase involved in protein sorting at the trans-Golgi network. The encoded protein may also act as a scaffold for the c-Jun N-terminal kinase signaling pathway, facilitating the formation of a functional signaling module. There are two named isoforms.

Gene ID:

57630

Application Details

Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid

Concentration: 1 μg/μL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Precaution of Use:

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months