

Datasheet for ABIN1419178

anti-Repetin antibody (AA 30-80) (HRP)[Go to Product page](#)

Overview

| | |
|----------------------|--|
| Quantity: | 100 µL |
| Target: | Repetin (RPTN) |
| Binding Specificity: | AA 30-80 |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This Repetin antibody is conjugated to HRP |
| Application: | Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)) |

Product Details

| | |
|-------------------|---|
| Immunogen: | KLH conjugated synthetic peptide derived from human Repetin |
| Isotype: | IgG |
| Cross-Reactivity: | Human, Mouse, Rat |
| Purification: | Purified by Protein A. |

Target Details

| | |
|-------------------|--|
| Target: | Repetin (RPTN) |
| Alternative Name: | Repetin (RPTN Products) |
| Background: | Synonyms: Intermediate filament associated protein, RPTN, RPTN_HUMAN. Background: Repetin is a 784 amino acid secreted protein that belongs to the S100-fused |

Target Details

protein family. Repetin is an extracellular epidermal matrix protein that is expressed in the epidermis and at high levels in eccrine sweat glands, the inner sheaths of hair roots and the filiform papilli of the tongue. A multifunctional epidermal matrix protein, repetin is involved in the cornified cell envelope formation. Containing two EF-hand domains, repetin's recombinant N-terminal domain binds calcium reversibly. Repetin and trichohyalin may compensate for absent loricrin. The repetin gene contains three exons, two introns and spans at least 5.6 kb. The repetin gene is conserved in cow, mouse and rat, and maps to human chromosome 1q21.3.

Gene ID: 126638

UniProt: [Q6XPR3](#)

Application Details

Application Notes: WB 1:300-5000
IHC-P 1:200-400

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling Advice: Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months