

[Go to Product page](#)

## Datasheet for ABIN1419640 **anti-RNF165 antibody (HRP)**

### Overview

Quantity:	100 µL
Target:	RNF165
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RNF165 antibody is conjugated to HRP
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human RNF165
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

### Target Details

Target:	RNF165
Alternative Name:	RNF165 ( <a href="#">RNF165 Products</a> )
Background:	<p>Synonyms: ring finger protein 165, RN165_HUMAN.</p> <p>Background: The RING-type zinc finger motif is present in a number of viral and eukaryotic proteins and is made of a conserved cysteine-rich domain that is able to bind two zinc atoms. Proteins that contain this conserved domain are generally involved in protein-protein</p>

## Target Details

interactions and protein-DNA interactions. Specifically, they are thought to be involved in the ubiquitination pathway of protein degradation. RNF165 (RING finger protein 165) is a 346 amino acid protein that contains one RING-type zinc finger. Via its RING-type zinc finger, RNF165 may play a role in transcriptional regulation and protein degradation events. The gene encoding RNF165 maps to human chromosome 18, which houses over 300 genes, including Trisomy 18, or Edwards syndrome, the second most common trisomy after Downs syndrome. Translocation between chromosome 18 and 14 is the most common translocation in cancers, and occurs in follicular lymphomas. Niemann-Pick disease, hereditary hemorrhagic telangiectasia and erythropoietic protoporphyria are also associated with chromosome 18.

Gene ID: 494470

## Application Details

Application Notes: WB 1:300-5000  
IHC-P 1:200-400

Restrictions: For Research Use only

## Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling Advice: Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months