

Datasheet for ABIN1420324
anti-CAF1p55 antibody (HRP)



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	CAF1p55 (CAF1)
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CAF1p55 antibody is conjugated to HRP
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CAF1p55
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

Target Details

Target:	CAF1p55 (CAF1)
Alternative Name:	CAF1p55 (CAF1 Products)
Background:	Synonyms: Caf 1, CAF-1, CAF-1 P55 subunit, CAF1 p55 subunit, CAF1A, CAF1P55, chaf1b, Chromatin assembly factor 1 p55 subunit, Chromatin assembly factor 1 p60 subunit, chromatin assembly factor 1 subunit, Chromatin assembly factor I, dCAF-1, dCAF1, Human chromatin assembly factor-I p55 subunit, Nucleosome remodeling factor 55 kDa subunit, NURF 55, NURF-

Target Details

55, p55, Probable histone binding protein Caf1, VPRBP_HUMAN.

Background: P55 is an extensively palmitoylated erythrocyte membrane protein, and a member of the MAGUK family. P55 also resists salt extraction, resulting in a high affinity for the plasma membrane. P55 contains a PDZ/DHR domain, a conserved SH-3 domain that appears to suppress tyrosine kinase activity of various oncoproteins, a 39-amino acid motif that binds to cytoskeletal protein 4.1R, and a guanylate kinase-like domain. Interaction with glycophorin C (GPC) and 4.1R suggests that p55 may play a role in the dynamic regulation in the erythrocyte membrane. In addition, p55 gene expression in vivo may be associated with a CpG island. P55 is constitutively expressed in K562 erythroleukemia cells during erythropoiesis and undergoes a 2-fold amplification after induction.

Gene ID: 9730

Application Details

Application Notes: WB 1:300-5000
IHC-P 1:200-400

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling Advice: Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months