antibodies .- online.com







anti-PDE4DIP antibody (HRP)



\sim					
()	VE	۲۱	/1	\triangle	Λ

Quantity:	100 μL	
Target:	PDE4DIP	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This PDE4DIP antibody is conjugated to HRP	
Application:	Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CMYA2/PDE4DIP	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Purification:	Purified by Protein A.	

Target Details

Target:	PDE4DIP	
Alternative Name:	CMYA2 (PDE4DIP Products)	
Background:	Synonyms: Cardiomyopathy associated protein 2, Cardiomyopathy-associated protein 2, CMYA2, MMGL, MYOME_HUMAN, Myomegalin, Pde4dip, Phosphodiesterase 4D interacting protein, Phosphodiesterase 4D-interacting protein.	
	Background: May function as an anchor sequestering components of the cAMP-dependent	

pathway to Golgi and/or centrosomes. Myomegalin, is a 2,346 amino acid protein that contains one NBPF domain and localizes to the nucleus, cytoplasm, centrosome and Golgi apparatus. Expressed at high levels in fetal and adult heart and at lower levels in brain and placenta, myomegalin is thought to function as an anchoring protein that sequesters members of the cAMP-dependent pathway to the Golgi and to centrosomes, thereby mediating cAMP pathway dynamics. Translocations in the gene that encodes myomegalin are associated with myeloproliferative disorders (MBDs), a group of diseases caused by an overproduction of blood cells. Myomegalin exists as twelve isoforms due to alternative splicing events.

Gene ID:

9659

Pathways:

SARS-CoV-2 Protein Interactome

Application Details

Application Notes:	IHC-P 1:200-400
Restrictions:	For Research Use only

Handling

- I arialing		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	