

[Go to Product page](#)

Datasheet for ABIN1421126

anti-C6orf192 antibody (AA 50-80) (Cy7)

Overview

| | |
|----------------------|---|
| Quantity: | 100 µL |
| Target: | C6orf192 |
| Binding Specificity: | AA 50-80 |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This C6orf192 antibody is conjugated to Cy7 |
| Application: | Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Western Blotting (WB) |

Product Details

| | |
|-------------------|--|
| Immunogen: | KLH conjugated synthetic peptide derived from human C6orf192 |
| Isotype: | IgG |
| Cross-Reactivity: | Human, Mouse, Rat |
| Purification: | Purified by Protein A. |

Target Details

| | |
|-------------------|--|
| Target: | C6orf192 |
| Alternative Name: | C6orf192 (C6orf192 Products) |
| Background: | Synonyms: C6orf192, dJ55C23.6, MFS-type transporter SLC18B1, Solute carrier family 18 member B1, SLC18B1 |

Target Details

Background: Making up nearly 6 % of the human genome, chromosome 6 contains around 1,200 genes within 170 million base pairs of sequence. Deletion of a portion of the q arm of chromosome 6 is associated with early onset intestinal cancer suggesting the presence of a cancer susceptibility locus. Porphyria cutanea tarda is associated with chromosome 6 through the HFE gene which, when mutated, predisposes an individual to developing this porphyria. Notably, the PARK2 gene, which is associated with Parkinson's disease, and the genes encoding the major histocompatibility complex proteins, which are key molecular components of the immune system and determine predisposition to rheumatic diseases, are also located on chromosome 6. Stickler syndrome, 21-hydroxylase deficiency and maple syrup urine disease are also associated with genes on chromosome 6. A bipolar disorder susceptibility locus has been identified on the q arm of chromosome 6. The C6orf192 gene product has been provisionally designated C6orf192 pending further characterization.

Gene ID: 116843

UniProt: [Q6NT16](#)

Application Details

Application Notes: IF(IHC-P) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months