# antibodies .- online.com





## anti-BEND2 antibody (HRP)



( )	11/	IN	/ie	A .
	/ // <del> </del>	۱ ات	/   (−	' \/\/

Quantity:	100 μL	
Target:	BEND2	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This BEND2 antibody is conjugated to HRP	
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human BEND2/CXorf56	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	Purified by Protein A.	

#### Target Details

rarget Details	
Target:	BEND2
Alternative Name:	BEND2 (BEND2 Products)
Background:	Synonyms: BEN domain-containing protein 2, BEN domain containing 2, BEND 2, BEND2, BEND-
	2, Chromosome X open reading frame 20, MGC33653, BEND2_HUMAN, CXorf56.

as two alternatively spliced isoforms and is considered a complete proteome. BEN domain

Background: BEND2 is a 799 amino acid protein that contains two BEN domains. BEND2 exists

mediates proteinDNA and proteinprotein interactions during chromatin organization and transcription. BEN domain may play a role in organization of viral DNA during replication or transcription. The BEND2 gene maps to human chromosome Xp22.13. Chromosome X consists of about 153 million base pairs and nearly 1,000 genes. The X and Y chromosomes are the human sex chromosomes. The combination of an X and Y chromosome lead to normal male development while two copies of X lead to normal female development. Color blindness, hemophilia, and Duchenne muscular dystrophy are well known X chromosome-linked conditions which affect males more frequently as males carry a single X chromosome.

### **Application Details**

Restrictions:	IHC-P 1:200-400  For Research Use only
Application Notes:	WB 1:300-5000

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months