antibodies.com

Datasheet for ABIN1421383 anti-SHARPIN antibody (AA 260-310) (Cy5.5)



Overview

| | 100 / |
|----------------------|---|
| Quantity: | 100 μL |
| Target: | SHARPIN |
| Binding Specificity: | AA 260-310 |
| Reactivity: | Human, Rat, Mouse |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This SHARPIN antibody is conjugated to Cy5.5 |
| Application: | Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human SHARPIN |
|-------------------|---|
| Isotype: | lgG |
| Cross-Reactivity: | Human, Mouse, Rat |
| Purification: | Purified by Protein A. |

Target Details

| Target: | SHARPIN |
|-------------------|--|
| Alternative Name: | SHARPIN (SHARPIN Products) |
| Background: | Synonyms: SIPL1, Sharpin, Shank-associated RH domain-interacting protein, Shank-interacting protein-like 1, hSIPL1, PSEC0216 |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN1421383 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

| | Background: Component of the LUBAC complex which conjugates linear polyubiquitin chains in |
|---------------------|---|
| | a head-to-tail manner to substrates and plays a key role in NF-kappa-B activation and regulation |
| | of inflammation. LUBAC conjugates linear polyubiquitin to IKBKG and RIPK1 and is involved in |
| | activation of the canonical NF-kappa-B and the JNK signaling pathways. Linear ubiquitination |
| | mediated by the LUBAC complex interferes with TNF-induced cell death and thereby prevents |
| | inflammation. LUBAC is proposed to be recruited to the TNF-R1 signaling complex (TNF-RSC) |
| | following polyubiquitination of TNF-RSC components by BIRC2 and/or BIRC3 and to conjugate |
| | linear polyubiquitin to IKBKG and possibly other components contributing to the stability of the |
| | complex. Together with FAM105B/otulin, the LUBAC complex regulates the canonical Wnt |
| | signaling during angiogenesis. |
| Gene ID: | 81858 |
| UniProt: | Q9H0F6 |
| Application Details | |
| Application Notes: | IF(IHC-P) 1:50-200 |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be |
| | handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |
| | |