# antibodies .- online.com





# anti-C16orf57 antibody (Cy5.5)



( )	ve	K\ /		A .
	$\cup$	1 V/	Щ.	V۷

Immunogen:	KLH conjugated synthetic pentide derived from human C16orf57	
Product Details		
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	
Conjugate:	This C16orf57 antibody is conjugated to Cy5.5	
Clonality:	Polyclonal	
Host:	Rabbit	
Reactivity:	Human, Mouse, Rat	
Target:	C16orf57 (USB1)	
Quantity:	100 μL	
Overview		

Immunogen:	KLH conjugated synthetic peptide derived from human C16orf57	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Purification:	Purified by Protein A.	

## **Target Details**

Target:	C16orf57 (USB1)	
Alternative Name:	C16orf57 (USB1 Products)	
Background:	Synonyms: Chromosome 16 open reading frame 57, CP057_HUMAN, FLJ13154, UPF0406 protein C16orf57.  Background: Involvement in disease, Defects in C16orf57 are the cause of poikiloderma with neutropenia (PN). PN is a genodermatosis characterized by poikiloderma, pachyonychia and	

#### **Target Details**

chronic neutropenia. The disorder starts as a papular erythematous rash on the limbs during the first year of life. It gradually spreads centripetally and, as the papular rash resolves, hypoand hyperpigmentation result, with development of telangiectasias. Another skin manifestation is pachyonychia, but alopecia and leukoplakia are distinctively absent. One of the most important extracutaneous symptoms is an increased susceptibility to infections, mainly affecting the respiratory system, primarily due to a chronic neutropenia and to neutrophil functional defects. Bone marrow abnormalities account for neutropenia and may evolve into myelodysplasia associated with the risk of leukemic transformation. Poikiloderma with neutropenia shows phenotypic overlap with Rothmund-Thomson syndrome.

Gene ID:

79650

### **Application Details**

Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months