antibodies .- online.com





anti-C16orf57 antibody (Cy7)



| \sim | | | | |
|--------|-----------|------|----|---|
| | $ V \cap$ | r\/I | 19 | ٨ |

| Quantity: | 100 μL | |
|--|---|--|
| Target: | C16orf57 (USB1) | |
| Reactivity: | Human, Mouse, Rat | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This C16orf57 antibody is conjugated to Cy7 | |
| Application: | Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |
| Product Details | | |
| | | |
| Immunogen: | KLH conjugated synthetic peptide derived from human C16orf57 | |
| Immunogen: Isotype: | KLH conjugated synthetic peptide derived from human C16orf57 | |
| | | |
| Isotype: | IgG | |
| Isotype: Cross-Reactivity: | IgG Human, Mouse, Rat | |
| Isotype: Cross-Reactivity: Purification: | IgG Human, Mouse, Rat | |
| Isotype: Cross-Reactivity: Purification: Target Details | lgG Human, Mouse, Rat Purified by Protein A. | |

Background: Involvement in disease, Defects in C16orf57 are the cause of poikiloderma with

neutropenia (PN). PN is a genodermatosis characterized by poikiloderma, pachyonychia and

protein C16orf57.

Target Details

chronic neutropenia. The disorder starts as a papular erythematous rash on the limbs during the first year of life. It gradually spreads centripetally and, as the papular rash resolves, hypoand hyperpigmentation result, with development of telangiectasias. Another skin manifestation is pachyonychia, but alopecia and leukoplakia are distinctively absent. One of the most important extracutaneous symptoms is an increased susceptibility to infections, mainly affecting the respiratory system, primarily due to a chronic neutropenia and to neutrophil functional defects. Bone marrow abnormalities account for neutropenia and may evolve into myelodysplasia associated with the risk of leukemic transformation. Poikiloderma with neutropenia shows phenotypic overlap with Rothmund-Thomson syndrome.

Gene ID:

79650

Application Details

| Application Notes: | IF(IHC-P) 1:50-200 | |
|--------------------|--|--|
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | 1 μg/μL | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. | |
| Storage: | -20 °C | |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | |
| Expiry Date: | 12 months | |