antibodies -online.com





anti-C16orf57 antibody (HRP)



Go to	D	1	
	Pron	וויאו וו	nane

()	1/0	r\ /1	014	
()	ve	I V I	-v	V

- OVERVIEW	
Quantity:	100 μL
Target:	C16orf57 (USB1)
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This C16orf57 antibody is conjugated to HRP
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human C16orf57
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.
Target Details	
Target:	C16orf57 (USB1)
Alternative Name:	C16orf57 (USB1 Products)
Background:	Synonyms: Chromosome 16 open reading frame 57, CP057_HUMAN, FLJ13154, UPF0406 protein C16orf57. Background: Involvement in disease, Defects in C16orf57 are the cause of poikiloderma with

neutropenia (PN). PN is a genodermatosis characterized by poikiloderma, pachyonychia and

chronic neutropenia. The disorder starts as a papular erythematous rash on the limbs during the first year of life. It gradually spreads centripetally and, as the papular rash resolves, hypoand hyperpigmentation result, with development of telangiectasias. Another skin manifestation is pachyonychia, but alopecia and leukoplakia are distinctively absent. One of the most important extracutaneous symptoms is an increased susceptibility to infections, mainly affecting the respiratory system, primarily due to a chronic neutropenia and to neutrophil functional defects. Bone marrow abnormalities account for neutropenia and may evolve into myelodysplasia associated with the risk of leukemic transformation. Poikiloderma with neutropenia shows phenotypic overlap with Rothmund-Thomson syndrome.

Gene ID:

79650

For Research Use only

Application Details

Application Notes:	WB 1:300-5000
	IHC-P 1:200-400

Handling

Restrictions:

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months