# antibodies -online.com





# anti-PCGF2 antibody (AA 170-210) (Cy5.5)



( )	11/	IN	/ie	A .
	/ // <del> </del>	۱ ات	/   (−	' \/\/

Quantity:	100 μL
Target:	PCGF2
Binding Specificity:	AA 170-210
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PCGF2 antibody is conjugated to Cy5.5
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human MEL18	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Purification:	Purified by Protein A.	

# Target Details

Target:	PCGF2
Alternative Name:	MEL18 (PCGF2 Products)
Background:	Synonyms: MEL-18, RNF110, ZNF144, Polycomb group RING finger protein 2, DNA-binding protein Mel-18, RING finger protein 110, Zinc finger protein 144, PCGF2, MEL18

Background: Transcriptional repressor. Binds specifically to the DNA sequence 5'-GACTNGACT-3'. Has tumor suppressor activity. May play a role in control of cell proliferation and/or neural cell development. Regulates proliferation of early T progenitor cells by maintaining expression of HES1. Also plays a role in antero-posterior specification of the axial skeleton and negative regulation of the self-renewal activity of hematopoietic stem cells. Component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1 complex acts via chromatin remodeling and modification of histones, it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility. Is not functionally redundant with BMI1, unlike BMI1 does not stimulate the E3 ubiquitin-protein ligase activity in a reconstituted PRC1-like complex (By similarity).

Gene ID: 7703

UniProt: P35227

Pathways: Chromatin Binding

## **Application Details**

Application Notes: IF(IHC-P) 1:50-200

Restrictions: For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months