antibodies -online.com





anti-GID8/C20orf11 antibody (Cy3)



\sim					
()	VE	۲۱	/1	\triangle	Λ

100 μL GID8/C20orf11 (GID8) Human, Mouse, Rat Rabbit Polyclonal This GID8/C20orf11 antibody is conjugated to Cy3 Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IE (p))	
Human, Mouse, Rat Rabbit Polyclonal This GID8/C20orf11 antibody is conjugated to Cy3	
Rabbit Polyclonal This GID8/C20orf11 antibody is conjugated to Cy3	
Polyclonal This GID8/C20orf11 antibody is conjugated to Cy3	
This GID8/C20orf11 antibody is conjugated to Cy3	
Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	
Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	
KLH conjugated synthetic peptide derived from human TWA1/C20orf11	
IgG	
Human, Mouse, Rat	
Purified by Protein A.	
GID8/C20orf11 (GID8)	
C20orf11 (GID8 Products)	
Synonyms: Chromosome 20 open reading frame 11, CT011_HUMAN, Protein C20orf11, Twa1,	
Two hybrid associated protein 1, Two hybrid-associated protein 1 with Ran BP-M, Two hybrid associated protein 1 with RanBPM, Two hybrid-associated protein 1 with RanBPM.	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn | International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com | Page 1/2 | Product datasheet for ABIN1421969 | 03/06/2024 | Copyright antibodies-online. All rights reserved.

Background: TWA1 is a 228 amino acid nuclear protein that is highly conserved throughout

evolution. TWA1 interacts with Ran BP-M, a protein that is implicated in a diverse array of cellular processes including DNA replication, entry into and exit from mitosis, and the transport of RNA and proteins through the nuclear pore complex. Together, Ran BP-M and TWA1 form a complex with cytoplasmic Muskelin, a mediator of cell spreading. It is thought that this complex plays a role in the Ran GTPase cycle and therefore a potential role in the cell cycle. TWA1 contains a LisH-CTLH motif which is usually found in proteins that are involved in nucleokinesis, chromosome segregation, cell migration and microtubule dynamics. The gene encoding TWA1 maps to human chromosome 20, which comprises approximately 2 % of the human genome and contains nearly 63 million bases that encode over 600 genes.

Application Details

Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	