

Datasheet for ABIN1423210

anti-CLEC11A antibody (HRP)



Go to Product page

_						
	V	\triangle	r۱	/1	\triangle	Λ/
	' V '		ΙV			v v

Quantity:	100 μL	
Target:	CLEC11A	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CLEC11A antibody is conjugated to HRP	
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human Pleckstrin/p47	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Purification:	Purified by Protein A.	
Target Details		
Target:	CLEC11A	
Alternative Name:	p47 (CLEC11A Products)	
Background:	Synonyms: p47, Platelet 47 kDa protein, Platelet and leukocyte C kinase substrate, Platelet P47,	
	Platelet p47 protein, Pleckstrin, PLEK, PLEK_HUMAN, FLJ27168, OTTHUMP00000159962.	
	Background: Platelet P47, also known as NSFL1C, UBX1, UBXD10 or UBXN2C, is a 370 amino	

acid protein that localizes to both the nucleus and the golgi apparatus (specifically to golgi

stacks) and contains one SEP domain and one UBX domain. Functioning as part of a ternary complex with VCP (a protein involved in the heterotypic fusion of transport vesicles with their target membranes) and Syntaxin 5, p47 interacts with and reduces the ATPase activity of VCP and is required for the fragmentation of golgi stacks during mitosis and for subsequent reassembly of golgi stacks after mitosis. p47 is subject to phosphorylation during mitosis, which inhibits p47-golgi interaction and is, therefore, required for proper golgi stack formation and cisternal regrowth. Human p47 shares 89 % sequence identity with its mouse counterpart, suggesting a conserved role between species. Multiple isoforms of p47 exist due to alternative splicing events.

Gene ID:

5341

For Research Use only

Application Details

Application Notes:	WB 1:300-5000
	IHC-P 1:200-400

Handling

Restrictions:

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months