antibodies -online.com





Datasheet for ABIN1423235

anti-CPN2 antibody (Cy3)



()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL	
Target:	CPN2	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CPN2 antibody is conjugated to Cy3	
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human CPN2	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Purification:	Purified by Protein A.	
Target Details		
Target:	CPN2	
Alternative Name:	CPN2 (CPN2 Products)	
Background:	Synonyms: CPN 2, CPN2, CPN-2, ACBP, Carboxypeptidase N 83 kDa chain, Carboxypeptidase N	
	large subunit, Carboxypeptidase N regulatory subunit, Carboxypeptidase N subunit 2	

[Precursor], carboxypeptidase N, polypeptide 2, CPN2_HUMAN.

Background: CPN2 is a zinc metalloprotease, and cleaves carboxy-terminal arginines and

Target Details

lysines from peptides found in the bloodstream such as complement anaphylatoxins, kinins, and creatine kinase MM (CK-MM). By removing only one amino acid, CPN has the ability to change peptide activity and receptor binding. It is a 280 kDa tetrameric glycoprotein that is synthesized by the liver and secreted into the plasma. It consists of 2 identical 83 kDa regulatory subunits (CPN2) and 2 identical 50 kDa catalytic subunits (CPN1). CPN2, the 83 kDa subunit, binds and stabilizes the catalytic subunit at 37 degrees Celsius and keeps it in circulation. Under some circumstances it may be an allosteric modifier of the catalytic subunit. CPN is a member of a larger family of carboxypeptidases, many of which also cleave arginine and lysine. Because of the highly conserved active sites and the possible redundant functions of carboxypeptidases, it has been difficult to elucidate the role of CPN in disease processes.

Gene ID:

1370

IF(IHC-P) 1:50-200

Application Details

Application Notes:

Application Notes.	IF(INC-P) 1.30-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	