.-online.com antibodies

## Datasheet for ABIN1423238 anti-CPN2 antibody (Cy7)



Overview	
Quantity:	100 μL
Target:	CPN2
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CPN2 antibody is conjugated to Cy7
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
	1/111 conjugated synthetic nontide deviced frame hymoen ODNO

Immunogen:	KLH conjugated synthetic peptide derived from human CPN2
Isotype:	lgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

## Target Details

Target:	CPN2
Alternative Name:	CPN2 (CPN2 Products)
Background:	Synonyms: CPN 2, CPN2, CPN-2, ACBP, Carboxypeptidase N 83 kDa chain, Carboxypeptidase N large subunit, Carboxypeptidase N regulatory subunit, Carboxypeptidase N subunit 2
	[Precursor], carboxypeptidase N, polypeptide 2, CPN2_HUMAN.
	Background: CPN2 is a zinc metalloprotease, and cleaves carboxy-terminal arginines and

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN1423238 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

	lysines from peptides found in the bloodstream such as complement anaphylatoxins, kinins,
	and creatine kinase MM (CK-MM). By removing only one amino acid, CPN has the ability to
	change peptide activity and receptor binding. It is a 280 kDa tetrameric glycoprotein that is
	synthesized by the liver and secreted into the plasma. It consists of 2 identical 83 kDa
	regulatory subunits (CPN2) and 2 identical 50 kDa catalytic subunits (CPN1). CPN2, the 83 kDa
	subunit, binds and stabilizes the catalytic subunit at 37 degrees Celsius and keeps it in
	circulation. Under some circumstances it may be an allosteric modifier of the catalytic subunit.
	CPN is a member of a larger family of carboxypeptidases, many of which also cleave arginine
	and lysine. Because of the highly conserved active sites and the possible redundant functions
	of carboxypeptidases, it has been difficult to elucidate the role of CPN in disease processes.
Gene ID:	1370
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months