antibodies .- online.com





anti-CWH43 antibody (AA 540-590) (HRP)



Go to	D	1	
	Proc	ויאו וו	Dane

/ N	11/0	K /	\sim	A /
	ve	1 \/	ı⊢۱	Λ
\sim	' V C	1 V		v v

Quantity:	100 μL
Target:	CWH43
Binding Specificity:	AA 540-590
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CWH43 antibody is conjugated to HRP
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CWH43
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

Target Details

Target:	CWH43
Alternative Name:	CWH43 (CWH43 Products)
Background:	Synonyms: Cell wall biogenesis protein 43 C-terminal homolog, Cwh43, imds 60,
	PG2IP_HUMAN, PGAP2-interacting protein, CWH43-C, PGAP2IP

Background: Representing approximately 6 % of the human genome, chromosome 4 contains nearly 900 genes. Notably, the Huntingtin gene, which is found to encode an expanded glutamine tract in cases of Huntington's disease, is on chromosome 4. FGFR-3 is also encoded on chromosome 4 and has been associated with thanatophoric dwarfism, achondroplasia, Muenke syndrome and bladder cancer. Chromosome 4 is also tied to Ellis-van Creveld syndrome, methylmalonic acidemia and polycystic kidney disease. Chromosome 4 reportedly contains the largest gene deserts (regions of the genome with no protein encoding genes) and has one of the two lowest recombination frequencies of the human chromosomes. The CWH43 gene product has been provisionally designated CWH43 pending further characterization.

Gene ID: 80157

UniProt: Q9H720

Pathways: Inositol Metabolic Process

Application Details

Application Notes: WB 1:300-5000

IHC-P 1:200-400

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months