

Datasheet for ABIN1429255

anti-NTSR2 antibody (PE)



| _ | | | | |
|----|-----|-----|-----|---|
| () | ve. | rv/ | 101 | Λ |

| Quantity: | 100 μL |
|--------------|---|
| Target: | NTSR2 |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This NTSR2 antibody is conjugated to PE |
| Application: | Western Blotting (WB) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human Neurotensin Receptor 2 | |
|-----------------------|--|--|
| Isotype: | IgG | |
| Cross-Reactivity: | Mouse | |
| Predicted Reactivity: | Human,Rat,Dog,Cow,Sheep,Rabbit | |
| Purification: | Purified by Protein A. | |

Target Details

| Target: | NTSR2 | |
|-------------------|---|--|
| Alternative Name: | Neurotensin Receptor 2 (NTSR2 Products) | |
| Background: | Synonyms: Levocabastine sensitive neurotensin receptor, Levocabastine-sensitive neurotensin | |
| | receptor, Neurotensin receptor type 2, NT R 2, NT-R-2, NTR2, NTR2 receptor, NTR2_HUMAN, | |

Ntsr2.

Background: Neurotensin (NT) initiates an intracellular response by interacting with the G protein-coupled receptors NTR1 (NTS1 receptor, high affinity NTR) and NTR2 (NTS2 receptor, levocabastine-sensitive neurotensin receptor), and the type I receptor NTR3 (NTS3 receptor, sortilin-1, Gp95). NT has a wide distribution in regions of the brain and in peripheral tissues where NT receptors can contribute to hypotension, hyperglycemia, hypothermia, antinociception and regulation of intestinal motility and secretion. HL-60 cells express NTR1, which can couple to Gq, Gi/o, or Gs. Alternative splicing of rat NTR2 can generate a 5-transmembrane domain variant isoform that is co-expressed with the full-length NTR2 throughout the brain and spinal cord. NTR3 activation in the murine microglial cell line N11 induces MIP-2, MCP-1, IL-1beta and TNF?in an ERK1/2 and Akt kinase-dependent manner.

Application Details

| Application Notes: | FCM(1:20-100) |
|--------------------|-----------------------|
| Restrictions: | For Research Use only |

Handling

| Format: | Liquid |
|--------------------|---|
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |