.-online.com antibodies

Datasheet for ABIN1437238 anti-IFI35 antibody (PE-Cy3)



Overview

Quantity:	100 µL
Target:	IFI35
Reactivity:	Human, Mouse, Rat, Cow, Dog, Pig
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IFI35 antibody is conjugated to PE-Cy3
Application:	Western Blotting (WB)
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human IFI35
lsotype:	lgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.
Target Details	
Target:	IFI35
Alternative Name:	IFI35 (IFI35 Products)
Background:	Synonyms: I 35, i-35, I35, P 35, P35, IN35, IN35_HUMAN, Interferon induced 35 kDa protein,
	Interferon-induced 35 kDa protein.
	Background: The Interferon family of proteins are able to alter the expression of a variety of
	target genes, thereby controlling various events within the cell. IFI-35 (Interferon-induced 35

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN1437238 | 03/06/2024 | Copyright antibodies-online. All rights reserved.

the nucleus and expressed in macrophages, fibroblasts and epithelial cells, IFI-35 is a l zipper protein that can form homodimers, but, unlike most leucine zipper proteins, can DNA. Upon induction by IFN-? IFI-35 associates with Nmi (N-Myc-interacting protein), r in the formation of a high molecular weight complex that is thought to play a role in IFI ?signaling and cellular responses. Once complexed with Nmi, IFI-35 is unable to be deer the proteasome, suggesting that IFI-35 is protected from degradation only when needer ?Two isoforms of IFI-35 exist due to alternative splicing events. Molecular Weight: 31kDa Gene ID: 3430 Application Details Application Notes: FCM(1:100-500) Optimal working dilution should be determined by the investigator. Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL. Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Procein 50 % Glycerol. Preservative: Sodium azide		
zipper protein that can form homodimers, but, unlike most leucine zipper proteins, can DNA. Upon induction by IFN-? IFI-35 associates with Nmi (N-Myc-interacting protein), t in the formation of a high molecular weight complex that is thought to play a role in IFI ?signaling and cellular responses. Once complexed with Nmi, IFI-35 is unable to be dependent to proteasome, suggesting that IFI-35 is protected from degradation only when needent the proteasome, suggesting that IFI-35 exist due to alternative splicing events. Molecular Weight: 31kDa Gene ID: 3430 Application Details FCM(1:100-500) Optimal working dilution should be determined by the investigator. Restrictions: For Research Use only Handling Iuquid Concentration: 1 µg/µL. Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclisto 90 % Glycerol. Preservative: Sodium azide Precaution of Use: This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE with should be handled by trained staff only.		kDa protein), also known as IFP35, is a 286 amino acid interferon-induced protein. Localized to
DNA. Upon induction by IFN-? IFI-35 associates with Nmi (N-Myc-interacting protein), r in the formation of a high molecular weight complex that is thought to play a role in IFI ?signaling and cellular responses. Once complexed with Nmi, IFI-35 is unable to be dependent of the proteasome, suggesting that IFI-35 is protected from degradation only when needent? ?Two isoforms of IFI-35 exist due to alternative splicing events. Molecular Weight: 31kDa Gene ID: 3430 Application Details Application Notes: FCM(1:100-500) Optimal working dilution should be determined by the investigator. Restrictions: For Research Use only Handling Iuquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclist 50 % Glycerol. Preservative: Sodium azide Precaution of Use: This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE witshould be handled by trained staff only.		the nucleus and expressed in macrophages, fibroblasts and epithelial cells, IFI-35 is a leucine
in the formation of a high molecular weight complex that is thought to play a role in IFI ?signaling and cellular responses. Once complexed with Nmi, IFI-35 is unable to be dependence. ?Two isoforms of IFI-35 exist due to alternative splicing events. Molecular Weight: 31kDa Gene ID: 3430 Application Details Application Notes: FCM(1:100-500) Optimal working dilution should be determined by the investigator. Restrictions: For Research Use only Handling Iuquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proceid 50 % Glycerol. Preservative: Sodium azide Precaution of Use: This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE w should be handled by trained staff only.		zipper protein that can form homodimers, but, unlike most leucine zipper proteins, cannot bind
?signaling and cellular responses. Once complexed with Nmi, IFI-35 is unable to be dependence. ?Two isoforms of IFI-35 exist due to alternative splicing events. Molecular Weight: 31kDa Gene ID: 3430 Application Details Application Notes: FCM(1:100-500) Optimal working dilution should be determined by the investigator. Restrictions: For Research Use only Handling Iuquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Prooling 50 % Glycerol. Preservative: Sodium azide Precaution of Use: This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE wishould be handled by trained staff only.		DNA. Upon induction by IFN-? IFI-35 associates with Nmi (N-Myc-interacting protein), resulting
the proteasome, suggesting that IFI-35 is protected from degradation only when needs ? Two isoforms of IFI-35 exist due to alternative splicing events. Molecular Weight: 31kDa Gene ID: 3430 Application Details Application Notes: FCM(1:100-500) Optimal working dilution should be determined by the investigator. Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proofing 50 % Glycerol. Preservative: Sodium azide Precaution of Use: This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE w should be handled by trained staff only.		in the formation of a high molecular weight complex that is thought to play a role in IFN-
? Two isoforms of IFI-35 exist due to alternative splicing events. Molecular Weight: 31kDa Gene ID: 3430 Application Details FCM(1:100-500) Optimal working dilution should be determined by the investigator. Optimal working dilution should be determined by the investigator. Restrictions: For Research Use only Handling Iuquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proofing 50 % Glycerol. Preservative: Sodium azide Precaution of Use: This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE with should be handled by trained staff only.		signaling and cellular responses. Once complexed with Nmi, IFI-35 is unable to be degraded by?
Molecular Weight: 31kDa Gene ID: 3430 Application Details Application Notes: FCM(1:100-500) Optimal working dilution should be determined by the investigator. Restrictions: For Research Use only Handling Induction Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proofing 50 % Glycerol. Preservative: Sodium azide Precaution of Use: This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE with the should be handled by trained staff only.		the proteasome, suggesting that IFI-35 is protected from degradation only when needed by IFN-
Gene ID: 3430 Application Details FCM(1:100-500) Application Notes: FCM(1:100-500) Optimal working dilution should be determined by the investigator. For Research Use only Handling For Research Use only Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Process Foreservative: Sodium azide Preservative: Sodium azide Precaution of Use: This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE w should be handled by trained staff only.		? Two isoforms of IFI-35 exist due to alternative splicing events.
Application Details FCM(1:100-500) Application Notes: FCM(1:100-500) Optimal working dilution should be determined by the investigator. Restrictions: Restrictions: For Research Use only Handling Iuquid Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Procting 50 % Glycerol. Preservative: Sodium azide Precaution of Use: This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE we should be handled by trained staff only.	Molecular Weight:	31kDa
Application Notes: FCM(1:100-500) Optimal working dilution should be determined by the investigator. Restrictions: For Research Use only Handling Iuquid Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Procling 50 % Glycerol. Preservative: Sodium azide Precaution of Use: This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE w should be handled by trained staff only.	Gene ID:	3430
Application Notes: FCM(1:100-500) Optimal working dilution should be determined by the investigator. Restrictions: For Research Use only Handling Iuquid Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Procling 50 % Glycerol. Preservative: Sodium azide Precaution of Use: This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE w should be handled by trained staff only.	Application Dataila	
Optimal working dilution should be determined by the investigator. Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Procling 50 % Glycerol. Preservative: Sodium azide Precaution of Use: This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE with a should be handled by trained staff only.	Application Details	
Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclision % Glycerol. Preservative: Sodium azide Precaution of Use: This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE with the should be handled by trained staff only.	Application Notes:	FCM(1:100-500)
Handling Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclision % Glycerol. Preservative: Sodium azide Precaution of Use: This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE with the should be handled by trained staff only.		Optimal working dilution should be determined by the investigator.
Format:LiquidConcentration:1 μg/μLBuffer:Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Procli 50 % Glycerol.Preservative:Sodium azidePrecaution of Use:This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE w should be handled by trained staff only.	Restrictions:	For Research Use only
Concentration:1 μg/μLBuffer:Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Procli 50 % Glycerol.Preservative:Sodium azidePrecaution of Use:This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE w should be handled by trained staff only.	Handling	
Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Procli 50 % Glycerol. Preservative: Sodium azide Precaution of Use: This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE w should be handled by trained staff only.	Format:	Liquid
50 % Glycerol. Preservative: Sodium azide Precaution of Use: This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE was should be handled by trained staff only.	Concentration:	1 μg/μL
Preservative: Sodium azide Precaution of Use: This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE was should be handled by trained staff only.	Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
Precaution of Use: This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE w should be handled by trained staff only.		50 % Glycerol.
should be handled by trained staff only.	Preservative:	Sodium azide
	Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
Storage: -20 °C		should be handled by trained staff only.
	Storage:	-20 °C
Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
	Expiry Date:	12 months