

Datasheet for ABIN1437250

anti-ARL6IP5 antibody (PE-Cy5.5)



Go to Product page

_				
	۱۱ / ۱	rv		۱۸/
	' V '	 ı v	Ι.	v v

Quantity:	100 μL	
Tanak	ADL CIDE (Adding)	
Target:	ARL6IP5 (Arl6ip5)	
Reactivity:	Human, Mouse, Rat, Cow, Dog, Pig	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This ARL6IP5 antibody is conjugated to PE-Cy5.5	
Application:	Western Blotting (WB)	

Product Details

lmmunogen:	KLH conjugated synthetic peptide derived from human JWA		
Isotype:	IgG		
Cross-Reactivity:	Human, Mouse, Rat		
Purification:	Purified by Protein A.		

Target Details

Target:	ARL6IP5 (Arl6ip5)
Alternative Name:	JWA (Arl6ip5 Products)
Background:	Synonyms: Addicsin, ADP ribosylation like factor 6 interacting protein 5, Aip 5, Aip 5, ARL 6
	interacting protein 5, ARL6IP5 antibody Cytoskeleton related vitamin A responsive protein,
	Dermal papilla derived protein 11, DERP 11, DERP11, Glutamate transporter EEAC 1 associated
	protein, Glutamate transporter EEAC1 associated protein, GTRAP3 18, Hp 22, Hp22, HSPC 127,

HSPC127, JM 5, JM5, JMX, PRA 1 domain family 3, PRA 2, PRA1 domain family 3, PRA2, PRAF 3, PRAF3, Prenylated Rab acceptor protein 2, Protein JWa, Putative MAPK activating protein PM27, PRAF3_HUMAN.

Background: JWA is a four-transmembrane environmental responsive protein which binds to the CC chemokine recepor 5 (CCR5), a major co-receptor for human immunodeficiency virus (HIV). JWA is involved in environmental stress-responsive pathways in K562 cells, an erythroleukemia cell line derived from patients with chronic myeloid leukemia. Environmental stressors to K562 cells such as heat shock, a higher temperature than the ideal body temperature of the organism from which the cell line was derived, and oxidative stress, the production of oxygen-centered free radicals, regulate and increase the expres-sion of JWA. This response to environmental stressors suggests similiarity of JWA to heat shock protein 70 (HSP70), which is upregulated by heat stress and toxic chemicals.

Molecular Weight: 21kDa

Gene ID: 10550

Pathways: Dicarboxylic Acid Transport

Application Details

Application Notes: FCM(1:100-500)

Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

ш	Jand	lina
г	land	1111()
•	10110	9

Expiry Date:

12 months