antibodies .- online.com





anti-MAP1LC3A antibody (N-Term, pSer12)





Publication



Go to Product page

_					
U	١V	e	rv	le	V

Quantity:	0.4 mL
Target:	MAP1LC3A
Binding Specificity:	AA 1-30, N-Term, pSer12
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MAP1LC3A antibody is un-conjugated
Application:	Western Blotting (WB), Dot Blot (DB), Enzyme Immunoassay (EIA)
Product Details	
Immunogen:	KLH conjugated synthetic phosphopeptide between 1-30 amino acids surrounding Ser12 of Human LC3
Isotype:	lg Fraction
Cross-Reactivity (Details):	Species reactivity (expected):Mouse, Rat, Bovine. Species reactivity (tested):Human.
Purification:	Protein A Affinity Chromatography. Then, the antibody fraction is peptide affinity purified in a 2-step procedure with peptides. The antibody is eluted with high and low pH buffers and neutralized immediately, followed by dialysis against PBS

Target Details

Target: MAP1LC3A


Target Details

Alternative Name:	LC3 (MAP1LC3A Products)	
Background:	MAP1A and MAP1B are microtubule-associated proteins which mediate the physical	
	interactions between microtubules and components of the cytoskeleton. These proteins are	
	involved in formation of autophagosomal vacuoles (autophagosomes). MAP1A and MAP1B	
	each consist of a heavy chain subunit and multiple light chain subunits. MAP1LC3a is one of	
	the light chain subunits and can associate with either MAP1A or MAP1B. The precursor	
	molecule is cleaved by APG4B/ATG4B to form the cytosolic form, LC3-I. This is activated by	
	APG7L/ATG7, transferred to ATG3 and conjugated to phospholipid to form the membrane-	
	bound form, LC3-II. Macroautophagy is the major inducible pathway for the general turnover of	
	cytoplasmic constituents in eukaryotic cells, it is also responsible for the degradation of active	
	cytoplasmic enzymes and organelles during nutrient starvation. Macroautophagy involves the	
	formation of double-membrane bound autophagosomes which enclose the cytoplasmic	
	constituent targeted for degradation in a membrane bound structure, which then fuse with the	
	lysosome (or vacuole) releasing a single-membrane bound autophagic bodies which are then	
	degraded within the lysosome (or vacuole). Synonyms: Autophagy-related protein LC3 A,	
	Autophagy-related ubiquitin-like modifier LC3 A, LC3A, MAP1 light chain 3-like protein 1, MAP1	
	/ 1B light chain 3 A, MAP1A/1B light chain 3 A, MAP1A/MAP1B LC3 A, MAP1LC3A, Microtubule	
	associated protein 1 light chain 3 alpha, Microtubule-associated proteins 1A/1B light chain 3A	
Molecular Weight:	14272 Da	
NCBI Accession:	NM_032514	
Pathways:	Autophagy	
Application Details		
Application Notes:	Optimal working dilution should be determined by the investigator.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	PBS, 0.09 % (W/V) Sodium Azide	
Preservative:	Sodium azide	
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	

Handling


Handling Advice:	Avoid repeated freezing and thawing.
Storage:	4 °C/-20 °C
Storage Comment:	Store undiluted at 2-8 °C for one month or (in aliquots) at -20 °C for longer.
Publications	
Product cited in:	Su, Chao, Huang, Weng, Jeng, Lai: "Rab5 and class III phosphoinositide 3-kinase Vps34 are involved in hepatitis C virus NS4B-induced autophagy." in: Journal of virology , Vol. 85, Issue 20, pp. 10561-71, (2011) (PubMed).

Images



Western Blotting


Image 1. Something like SH-SY5Y cells expressing GFP-LC3-WT or-S12D treated with rapamycin or vehicle for 1h.



Dot Blot

Image 2. Dot blot analysis of Phospho-LC3 (APG8a) - Ser12 Antibody and Non phospho-LC3 (APG8a) Antibody on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are $0.5 \, \mu \text{g/ml}$.

Dot Blot



Western Blotting

Image 3. Immunoblots of phosphorylated LC3 (phospho-LC3) in CHO cell culture. LC3 and LC3 S12A mutant vectors were transfected into CHO cells. The cell lysates were separated with SDS-PAGE and blotted with anti-phospho-LC3 S12 antibody. LC3 = microtubule-associated protein light chain-3; S12A = replacement of the amino acid position 12 serine of LC3 with alanine. WT = wildtype LC3-transfected cell lysates; S12A = LC3 S12A mutant-transfected cell lysates; Empty vector = vector with no LC3 gene. Molecular size: LC3-I = 16kDa, and LC3-II = 14 kDa

Please check the product details page for more images. Overall 5 images are available for ABIN1449623.