

Datasheet for ABIN1449627
anti-LC3B antibody (C-Term)



[Go to Product page](#)

3 Images

Overview

Quantity:	0.4 mL
Target:	LC3B (MAP1LC3B)
Binding Specificity:	AA 88-118, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This LC3B antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Enzyme Immunoassay (EIA)

Product Details

Immunogen:	KLH conjugated synthetic peptide between 88~118 amino acids from the C-terminal region of Human LC3
Isotype:	Ig Fraction
Purification:	Saturated Ammonium Sulfate precipitation followed by dialysis against PBS

Target Details

Target:	LC3B (MAP1LC3B)
Alternative Name:	LC3B (MAP1LC3B Products)
Background:	Macroautophagy is the major inducible pathway for the general turnover of cytoplasmic constituents in eukaryotic cells, it is also responsible for the degradation of active cytoplasmic

Target Details

enzymes and organelles during nutrient starvation. Macroautophagy involves the formation of double-membrane bound autophagosomes which enclose the cytoplasmic constituent targeted for degradation in a membrane bound structure, which then fuse with the lysosome (or vacuole) releasing a single-membrane bound autophagic bodies which are then degraded within the lysosome (or vacuole). MAP1A and MAP1B are microtubule-associated proteins which mediate the physical interactions between microtubules and components of the cytoskeleton. These proteins are involved in formation of autophagosomal vacuoles (autophagosomes). MAP1A and MAP1B each consist of a heavy chain subunit and multiple light chain subunits. MAP1LC3b is one of the light chain subunits and can associate with either MAP1A or MAP1B. The precursor molecule is cleaved by APG4B/ATG4B to form the cytosolic form, LC3-I. This is activated by APG7L/ATG7, transferred to ATG3 and conjugated to phospholipid to form the membrane-bound form, LC3-II. Synonyms: Autophagy-related protein LC3 B, Autophagy-related ubiquitin-like modifier LC3 B, MAP1A / 1B light chain 3 B, MAP1A/MAP1B, MAP1LC3B, Map1alc3, Map1lc3, Map1lc3b, Microtubule-associated protein 1 light chain 3 beta, Microtubule-associated proteins 1A/1B light chain 3B

Gene ID:	81631
NCBI Accession:	NP_073729
Pathways:	Autophagy

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	0.25 mg/mL
Buffer:	PBS containing 0.09 % (W/V) Sodium Azide as preservative
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freezing and thawing.

Handling


Storage:	4 °C/-20 °C
Storage Comment:	Store undiluted at 2-8 °C for one month or (in aliquots) at -20 °C for longer.

Images




Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Formalin-Fixed, Paraffin-Embedded human cancer tissue reacted with LC3B Antibody (C-term) followed which was peroxidase-conjugated to the secondary antibody, followed by DAB staining.



Immunohistochemistry (Paraffin-embedded Sections)

Image 2. Formalin-Fixed, Paraffin-embedded human testis tissue reacted with Autophagy LC3B Antibody (C-term) followed which was peroxidase-conjugated to the secondary antibody, followed by DAB staining.



Western Blotting

Image 3. Western blot analysis of Autophagy LC3B Antibody (C-term) in 293 cell line lysates transiently transfected with the LC3 (APG8b) gene (2ug/lane).