

Datasheet for ABIN1449802

anti-ZNF265 antibody (Isoform 2, Middle Region)



[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	ZNF265 (Zranb2)
Binding Specificity:	Isoform 2, Middle Region
Reactivity:	Human, Mouse, Rat, Dog, Cow, Pig, Chicken, Xenopus laevis
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ZNF265 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	A synthetic peptide directed towards the middle region of human ZRANB2 (Isoform 2)
Sequence:	EDKESEGEEE DEDEDLSKYK LDEDEDEDDA DLSKYNLDAS EEEDSNKKKS
Cross-Reactivity (Details):	Species reactivity (tested): Human, Mouse, Rat, Dog, Bovine, Pig, Chicken, African clawed frog
Purification:	Purified using peptide immunoaffinity column

Target Details

Target:	ZNF265 (Zranb2)
Alternative Name:	ZRANB2 (Zranb2 Products)
Background:	ZRANB2 is a splice factor required for alternative splicing of SFRS10/TRA2B transcripts. It may interfere with constitutive 5'-splice site selection. Synonyms: ZIS, ZNF265, Zinc finger Ran-

Target Details

	binding domain-containing protein 2, Zinc finger protein 265, Zinc finger splicing
Molecular Weight:	36 kDa (320 aa)
Gene ID:	9406
NCBI Accession:	NP_005446


Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

Handling


Reconstitution:	Add 50 µL of distilled water to a final concentration of 1.0 mg/mL.
Concentration:	1.0 mg/mL after reconstitution
Handling Advice:	Avoid repeated freezing and thawing.
Storage:	4 °C/-20 °C
Storage Comment:	Store lyophilized at 2-8 °C or at -20 °C long term. After reconstitution store the antibody undiluted at 2-8 °C for up to one month or in aliquots at -20 °C long term.

Images



Immunohistochemistry

Image 1. Human Lung; ZRANB2 antibody - middle region in Human Lung cells using Immunohistochemistry



Western Blotting

Image 2. Human HepG2; WB Suggested Anti-ZRANB2 Antibody Titration: 0.2-1 ug/ml. Positive Control: HepG2 cell lysate; ZRANB2 antibody - middle region in Human HepG2 cells using Western Blot