

Datasheet for ABIN1449807

anti-ZNF92 antibody (C-Term, Isoform 2)[Go to Product page](#)**1** Image

Overview

Quantity:	0.1 mg
Target:	ZNF92
Binding Specificity:	C-Term, Isoform 2
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ZNF92 antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	A synthetic peptide directed towards the C-terminal region of human ZNF92 (isoform 2, Q03936-2)
Sequence:	KPYKYECDK AFNKFSTLIT HQIYTGEKP CKHECGRAFN KSSNYTKEKL
Cross-Reactivity (Details):	Species reactivity (tested): Human, Mouse
Purification:	Purified using Protein A affinity column

Target Details

Target:	ZNF92
Alternative Name:	ZNF92 (ZNF92 Products)
Background:	ZNF92, belonging to the krueppel C2H2-type zinc-finger protein family, is expressed early during

Target Details

	embryonic development and may be involved in transcriptional regulation.Synonyms: HTF12, Zinc finger protein 92, Zinc finger protein HTF12
Molecular Weight:	60 kDa (517 aa)
Gene ID:	168374
NCBI Accession:	NP_009070


Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

Handling

Reconstitution:	Add 100 µL of distilled water to a final concentration of 1 mg/mL.
Concentration:	1.0 mg/mL
Handling Advice:	Avoid repeated freezing and thawing.
Storage:	4 °C/-20 °C
Storage Comment:	Store lyophilized at 2-8 °C or at -20 °C long term. After reconstitution store the antibody undiluted at 2-8 °C for up to one month or in aliquots at -20 °C long term.

Images



Western Blotting

Image 1. Human Jurkat; WB Suggested Anti-ZNF92 Antibody Titration: 2.5ug/ml. ELISA Titer: 1:62500. Positive Control: Jurkat cell lysate; ZNF92 antibody - C-terminal region in Human Jurkat cells using Western Blot