antibodies -online.com





Datasheet for ABIN1493904

anti-TREML2 antibody (AA 188-237)

3 Images



Go to Product page

Overview

Quantity:	50 μg
Target:	TREML2
Binding Specificity:	AA 188-237
Reactivity:	Human, Monkey
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TREML2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunohistochemistry (Paraffinembedded Sections) (IHC (p))

Product Details

Brand:	IHC-plus™
Immunogen:	Synthetic peptide located between aa188-237 of human TREML2 (Q5T2D2, NP_079083). Percent identity by BLAST analysis: Human, Gorilla, Gibbon, Monkey (100%).
	refeetit identity by BE 161 dinaryole. Flamar, Comia, Classon, Monkey (100 kg.
	Type of Immunogen: Synthetic peptide
Isotype:	IgG
Specificity:	Human TREML2
Predicted Reactivity:	Percent identity by BLAST analysis: Human (100%).
Purification:	Immunoaffinity purified

Target Details

Target:	TREML2
Alternative Name:	TLT2 / TREML2 (TREML2 Products)
Background:	Name/Gene ID: TREML2
	Synonyms: TREML2, C6orf76, TLT2, DJ238023.1, TLT-2, Trem-like transcript 2 protein
Gene ID:	79865
NCBI Accession:	NP_079083
UniProt:	Q5T2D2

Application Details

Application Notes:	Approved: IHC, IHC-P (2.5 - 5 μg/mL), WB
	Usage: ELISA titer using peptide based assay: 1:62500. Western Blot: Suggested dilution at 1 μ g/mL in 5 % skim milk / PBS buffer, and HRP conjugated anti-Rabbit IgG should be diluted in 1:50000 - 100000 as second antibody.
Comment:	Target Species of Antibody: Human
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Distilled Water.
Concentration:	Lot specific
Buffer:	Lyophilized from PBS with 2 % sucrose
Handling Advice:	Avoid repeat freeze-thaw cycles.
Storage:	4 °C,-20 °C
Storage Comment:	Long term: -20°C, the use of 50% glycerol is recommended if storing aliquots in -20°C for long
	term use (up to 1 year)
	Short term (less than 1 week): 4°C. Avoid freeze-thaw cycles.




Image 1.





Image 2.



Immunohistochemistry

Image 3. Anti-TLT2 / TREML2 antibody IHC of human placenta. Immunohistochemistry of formalin-fixed, paraffinembedded tissue after heat-induced antigen retrieval. Antibody dilution 2.5-5 ug/ml. This image was taken for the unconjugated form of this product. ...