

Datasheet for ABIN1531316 anti-FAK antibody (pTyr407)

2 Images



Overview

Overview	
Quantity:	100 μL
Target:	FAK (PTK2)
Binding Specificity:	AA 373-422, pTyr407
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FAK antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC)
Product Details	
Immunogen:	The antiserum was produced against synthesized peptide derived from human FAK around the phosphorylation site of Tyr407.
Isotype:	IgG
Specificity:	FAK (Phospho-Tyr407) Antibody detects endogenous levels of FAK only when phosphorylated at Tyr407. PhosphorylationH:Y407 M:Y438 R:Y407
Purification:	The antibody was purified from rabbit antiserum by affinity-chromatography using phospho peptide. The antibody against non-phospho peptide was removed by chromatography using corresponding non-phospho peptide.
Purity:	> 95 %

Target Details


Target:	FAK (PTK2)
Alternative Name:	FAK (PTK2 Products)
Background:	Synonyms: FADK 1, FAK1, Focal adhesion kinase 1, pp125FAK, Protein- tyrosine kinase 2, PTK2 NCBI Gene Symbol: PTK2
Molecular Weight:	119 kDa
Gene ID:	5747
OMIM:	600758
OMIM: UniProt:	600758 Q05397

Application Details

Application Notes:	WB: 1:500~1:1000 IHC: 1:50~1:100 ELISA: 1:5000
Comment:	Unigene-Number: Hs.395482 (NCBI Gene Symbol: PTK2)
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Stable at -20°C for at least 1 year.
Expiry Date:	12 months



Immunohistochemistry

Image 1. Immunohistochemistry analysis of paraffinembedded human brain, using FAK (Phospho-Tyr407) Antibody. The picture on the right is treated with the synthesized peptide.



Western Blotting

Image 2. Western blot analysis of extracts from COS7 cells treated with EGF 200ng/ml 30', using FAK (Phospho-Tyr407) Antibody. The lane on the right is treated with the synthesized peptide.