

Datasheet for ABIN1532159
anti-NFATC4 antibody (pSer168)

2 Images

[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	NFATC4
Binding Specificity:	AA 136-185, pSer168
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NFATC4 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	The antiserum was produced against synthesized peptide derived from human NFAT3 around the phosphorylation site of Ser168 and Ser170.
Isotype:	IgG
Specificity:	NFAT3 (Phospho-Ser168+Ser170) Antibody detects endogenous levels of NFAT3 only when phosphorylated at Ser168 and Ser170. PhosphorylationH:S168+S170 M:S168+S170
Purification:	The antibody was purified from rabbit antiserum by affinity-chromatography using phospho peptide. The antibody against non-phospho peptide was removed by chromatography using corresponding non-phospho peptide.
Purity:	> 95 %

Target Details


Target:	NFATC4
Alternative Name:	NFAT3 (NFATC4 Products)
Background:	Synonyms: NF-AT3, NF-ATc4, NFAC4, NFATC4, NFC4, Nuclear factor of activated T-cells, cytoplasmic 4, T cell transcription factor NFAT3, Transcription complex subunit NF-ATc4 NCBI Gene Symbol: NFATC4
Molecular Weight:	95 kDa
Gene ID:	4776
OMIM:	602699
UniProt:	Q14934
Pathways:	RTK Signaling , WNT Signaling

Application Details

Application Notes:	IHC: 1:50~1:100 ELISA: 1:1000
Comment:	Unigene-Number: Hs.77810 (NCBI Gene Symbol: NFATC4)
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Stable at -20°C for at least 1 year.
Expiry Date:	12 months



Immunohistochemistry

Image 1. Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using NFAT3 (Phospho-Ser168+Ser170) Antibody. The picture on the right is treated with the synthesized peptide.



Western Blotting

Image 2.