

Datasheet for ABIN1532224

anti-Tyrosine Hydroxylase antibody (AA 41-90)



[Go to Product page](#)

3 Images

Overview

Quantity:	100 µL
Target:	Tyrosine Hydroxylase (TH)
Binding Specificity:	AA 41-90
Reactivity:	Human, Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Tyrosine Hydroxylase antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), ELISA

Product Details

Immunogen:	The antiserum was produced against synthesized peptide derived from human Tyrosine Hydroxylase.
Isotype:	IgG
Specificity:	Tyr Hydroxylase Antibody detects endogenous levels of total Tyr Hydroxylase protein.
Purification:	The antibody was purified from rabbit antiserum by affinity-chromatography using immunogen.
Purity:	> 95 %

Target Details

Target:	Tyrosine Hydroxylase (TH)
Alternative Name:	Tyrosine Hydroxylase (TH Products)

Target Details


Background:	Synonyms: TH isoform 3, TH isoform a, TH-4, TY3H, TYH, Tyrosine 3-hydroxylase, Tyrosine 3-monooxygenase, tyrosine hydroxylase NCBI Gene Symbol: TH
Molecular Weight:	58 kDa
Gene ID:	7054
OMIM:	191290
UniProt:	P07101
Pathways:	Dopaminergic Neurogenesis , Response to Water Deprivation , Sensory Perception of Sound , Carbohydrate Homeostasis , Feeding Behaviour

Application Details

Application Notes:	WB: 1:500~1:1000 IHC: 1:50~1:100 IF: 1:100~1:500 ELISA: 1:20000
Comment:	Unigene-Number: Hs.435609 (NCBI Gene Symbol: TH)
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Stable at -20°C for at least 1 year.
Expiry Date:	12 months




Western Blotting

Image 1. Western blot analysis of extracts from NIH-3T3 cells, treated with Forskolin 40nM 30', using Tyrosine Hydroxylase (Ab-40) Antibody. The lane on the right is treated with the synthesized peptide.



Immunohistochemistry

Image 2. Immunohistochemistry analysis of paraffin-embedded human brain tissue, using Tyrosine Hydroxylase (Ab-40) Antibody. The picture on the right is treated with the synthesized peptide.



Immunofluorescence

Image 3. Immunofluorescence analysis of HuvEc cells, using Tyrosine Hydroxylase (Ab-40) Antibody. The picture on the right is treated with the synthesized peptide.