

Datasheet for ABIN1532903

anti-NFKBIA antibody (AA 9-58)

2 Images



Go to Product page

\sim			
0^{\vee}	6	rVI	Ie.W

Quantity:	100 μL
Target:	NFKBIA
Binding Specificity:	AA 9-58
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NFKBIA antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC)
Product Details	
Immunogen:	The antiserum was produced against synthesized peptide derived from human IkappaB-alpha.
Isotype:	IgG
Specificity:	IkappaB-alpha Antibody detects endogenous levels of total IkappaB-alpha protein.
Purification:	The antibody was purified from rabbit antiserum by affinity-chromatography using immunogen.
Purity:	> 95 %
Target Details	
Target Details Target:	NFKBIA
	NFKBIA IkappaB-alpha (NFKBIA Products)


Target Details

	enhancer-binding protein MAD3, NF-kappaB inhibitor alpha, NFKBI, NFKBIA, RL/IF-1 NCBI Gene Symbol: NFKBIA
Molecular Weight:	35 kDa
Gene ID:	4792
OMIM:	164008
UniProt:	P25963
Pathways:	NF-kappaB Signaling, TCR Signaling, TLR Signaling, Fc-epsilon Receptor Signaling Pathway, Activation of Innate immune Response, Cellular Response to Molecule of Bacterial Origin, Maintenance of Protein Location, Hepatitis C, Protein targeting to Nucleus, Toll-Like Receptors Cascades, BCR Signaling

Application Details


Application Details	
Application Notes:	WB: 1:500~1:1000 IHC: 1:50~1:100 ELISA: 1:10000
Comment:	Unigene-Number: Hs.81328 (NCBI Gene Symbol: NFKBIA)
Restrictions:	For Research Use only
Handling	
Format:	Liquid

Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Stable at -20°C for at least 1 year.
Expiry Date:	12 months



Western Blotting

Image 1. Western blot analysis of extracts from HuvEc cells, treated with TNF-alpha 20ng/ml 30', using IkappaB-alpha (Ab-42) Antibody. The lane on the right is treated with the synthesized peptide.



Immunohistochemistry

Image 2. Immunohistochemistry analysis of paraffinembedded human breast carcinoma tissue, using IkappaBalpha (Ab-42) Antibody. The picture on the right is treated with the synthesized peptide.