antibodies -online.com





anti-PRKX antibody (AA 251-300)

2 Images



Go to Product page

Overview

| Quantity: | 100 μg |
|----------------------|--|
| Target: | PRKX |
| Binding Specificity: | AA 251-300 |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This PRKX antibody is un-conjugated |
| Application: | Western Blotting (WB), ELISA, Immunohistochemistry (IHC) |

Product Details

| Immunogen: | The antiserum was produced against synthesized peptide derived from human PRKX. |
|---------------|---|
| Isotype: | IgG |
| Specificity: | PRKX Antibody detects endogenous levels of total PRKX protein. |
| Purification: | The antibody was purified from rabbit antiserum by affinity-chromatography using immunogen. |
| Purity: | > 95 % |

Target Details

| Target: | PRKX |
|-------------------|---|
| Alternative Name: | PRKX (PRKX Products) |
| Background: | Synonyms: Serine/threonine-protein kinase PRKX, Protein kinase PKX1, PRKX |

Target Details


| | NCBI Gene Symbol: PRKX |
|-------------------|---------------------------|
| Molecular Weight: | 40 kDa |
| Gene ID: | 5613 |
| OMIM: | 300083 |
| UniProt: | P51817 |
| Pathways: | Thyroid Hormone Synthesis |

Application Details

| Application Notes: | WB: 1:500~1:1000 IHC: 1:50~1:100 ELISA: 1:20000 |
|--------------------|--|
| Comment: | Unigene-Number: Hs.390788 (NCBI Gene Symbol: PRKX) |
| Restrictions: | For Research Use only |


Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 mg/mL |
| Buffer: | phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol. |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Stable at -20°C for at least 1 year. |
| Expiry Date: | 12 months |



Western Blotting

Image 1. Western blot analysis of extracts from MCF-7 cells, using PRKX Antibody. The lane on the right is treated with the synthesized peptide.



Immunohistochemistry

Image 2. Immunohistochemistry analysis of paraffinembedded human breast carcinoma tissue, using PRKX Antibody. The picture on the right is treated with the synthesized peptide.