
antibodies -online.com







anti-TP53RK antibody (C-Term)

2 Images



Go to Product page

)\/(

Overview	
Quantity:	400 μL
Target:	TP53RK
Binding Specificity:	AA 221-250, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TP53RK antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	This TP53RK antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 221-250 amino acids from the C-terminal region of human TP53RK.
Clone:	RB36762
Isotype:	lg Fraction
Predicted Reactivity:	М
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.
Target Details	
Target:	TP53RK
Alternative Name:	TP53RK (TP53RK Products)

Target Details


Background:	Protein kinase that phosphorylates 'Ser-15' of p53/TP53 protein and may therefore participate in its activation.
Molecular Weight:	28160
Gene ID:	112858
NCBI Accession:	NP_291028
UniProt:	Q96S44

Application Details

Application Notes:	WB: 1:1000. WB: 1:1000
Restrictions:	For Research Use only


Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	TP53RK Antibody (C-term) can be refrigerated at 2-8 °C for up to 6 months. For long term storage, keep at -20 °C.
Expiry Date:	6 months



Western Blotting

Image 1. TP53RK Antibody (C-term) (ABIN1536869 and ABIN2848663) western blot analysis in HL-60 cell line lysates (35 μ g/lane).This demonstrates the TP53RK antibody detected the TP53RK protein (arrow).



Western Blotting

Image 2. All lanes: Anti-TP53RK Antibody (C-term) at 1:1000 dilution Lane 1: Jurkat whole cell lysate Lane 2: K562 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 28 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.